



C–H Functionalization/activation in organic synthesis

Richmond Sarpong

Editorial

Open Access

Address:
Department of Chemistry, University of California, Berkeley, 841A
Latimer Hall, Berkeley, CA 94720, USA

Email:
Richmond Sarpong - rsarpong@berkeley.edu

Keywords:
C–H activation; C–H functionalization

Beilstein J. Org. Chem. **2016**, *12*, 2315–2316.
doi:10.3762/bjoc.12.224

Received: 19 September 2016
Accepted: 29 September 2016
Published: 03 November 2016

This article is part of the Thematic Series "C–H
Functionalization/activation in organic synthesis".

Guest Editor: R. Sarpong

© 2016 Sarpong; licensee Beilstein-Institut.
License and terms: see end of document.

The last decade has seen an explosion in research reports in the area of C–H functionalization/activation in organic synthesis. This is not surprising. What is surprising is that it took many in the synthetic organic community this long to fully embrace this exciting and enabling technology. After all, organic compounds mainly consist of a carbon skeleton that bears a large number of hydrogens. It is therefore highly desirable to be able to take advantage of the myriad of C–H groups in organic molecules as functional handles for bond formation, and in some cases, bond-breaking processes.

“Modern” C–H functionalization/activation can trace its roots to the “clarion call” by Bergman and co-workers [1] in a famous *Accounts of Chemical Research* paper in 1995 where the potential for this reaction was discussed. The community has responded and today, there are investigators from all branches of chemistry, chemical biology, and engineering that are pursuing new powerful methods and strategies to achieve C–H functionalization/activation reactions. The growth in popularity of C–H functionalization/activation in organic synthesis can be attributed to the desire to implement more sustainable methods for synthesis and to achieve novel reactivity and selectivity in building molecules. A lot of exciting progress has been made in

a short period of time and the stage has been set for even more far-reaching developments in the future.

In this Thematic Series, a collection of 10 contributions from researchers in the area of C–H functionalization from Europe, United States, Japan, China, India, and Brazil is presented. These contributions include full accounts on primary research in the area of C–H functionalization/activation and reviews that focus on aspects of this exciting field. The powerful C_{sp^2} functionalization of heterocycles provides access to value-added compounds as reported by Itami et al. [2], Chatani et al. [3] and Waser et al. [4]. Alternatively, the C_{sp^2} functionalization of benzenoids is equally powerful and affords interesting opportunities as described by Lipshutz et al. [5], Bisai et al. [6] and Chen et al. [7]. The emerging power of controlled and selective C_{sp^3} functionalization is also captured in this series with contributions from Wang et al. [8], May et al. [9], Machado et al. [10] and Dai et al. [11].

I am grateful to all those that have participated in making this Thematic Series, first and foremost, the authors. I am also thankful to the reviewers, who made many constructive suggestions. Overall, it is clear that the field of C–H functionalization/

activation in all its varied forms is an exciting area of modern synthetic chemistry and in many ways is revolutionizing synthetic organic chemistry. This Thematic Series provides a view into this burgeoning and exciting world.

Richmond Sarpong

Berkeley, September 2016

References

1. Arndtsen, B. A.; Bergman, R. G.; Mobley, T. A.; Peterson, T. H. *Acc. Chem. Res.* **1995**, *28*, 154–162. doi:10.1021/ar00051a009
2. Hata, K.; Ito, H.; Segawa, Y.; Itami, K. *Beilstein J. Org. Chem.* **2015**, *11*, 2737–2746. doi:10.3762/bjoc.11.295
3. Tobisu, M.; Igarashi, T.; Chatani, N. *Beilstein J. Org. Chem.* **2016**, *12*, 654–661. doi:10.3762/bjoc.12.65
4. Tolnai, G. L.; Brand, J. P.; Waser, J. *Beilstein J. Org. Chem.* **2016**, *12*, 745–749. doi:10.3762/bjoc.12.74
5. Nishikata, T.; Abela, A. R.; Huang, S.; Lipshutz, B. H. *Beilstein J. Org. Chem.* **2016**, *12*, 1040–1064. doi:10.3762/bjoc.12.99
6. Kumar, N.; Ghosh, S.; Bhunia, S.; Bisai, A. *Beilstein J. Org. Chem.* **2016**, *12*, 1153–1169. doi:10.3762/bjoc.12.111
7. Nack, W. A.; Wang, X.; Wang, B.; He, G.; Chen, G. *Beilstein J. Org. Chem.* **2016**, *12*, 1243–1249. doi:10.3762/bjoc.12.119
8. Wang, B.; Qiu, D.; Zhang, Y.; Wang, J. *Beilstein J. Org. Chem.* **2016**, *12*, 796–804. doi:10.3762/bjoc.12.78
9. Shih, J.-L.; Chen, P.-A.; May, J. A. *Beilstein J. Org. Chem.* **2016**, *12*, 985–999. doi:10.3762/bjoc.12.97
10. Santiago, J. V.; Machado, A. H. L. *Beilstein J. Org. Chem.* **2016**, *12*, 882–902. doi:10.3762/bjoc.12.87
11. Ye, Z.; Gettys, K. E.; Dai, M. *Beilstein J. Org. Chem.* **2016**, *12*, 702–715. doi:10.3762/bjoc.12.70

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the *Beilstein Journal of Organic Chemistry* terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at:
[doi:10.3762/bjoc.12.224](https://doi.org/10.3762/bjoc.12.224)