

Supporting Information

for

Functionalization of anthracene: A selective route to brominated 1,4-anthraquinones

Kiyemet Berkil Akar¹, Osman Cakmak ^{*1}, Orhan Büyükgüngör^{2,§}, Ertan Sahin^{3,§}

Address: ¹Department of Chemistry, Faculty of Art and Science, Gaziosmanpasa University, 60250, Tokat, Turkey; ²Department of Physics, Faculty of Art and Science, Ondokuzmayıs University, 55060, Samsun, Turkey and ³Department of Chemistry, Faculty of Art and Science, Atatürk University, 25240, Erzurum, Turkey

Email: Kiyemet Berkil Akar - kiyemetberkil@gmail.com, *Osman Cakmak - cakmak.osman@gmail.com, Orhan Büyükgüngör - orhanb@omu.edu.tr, Ertan Sahin - ertan@atauni.edu.tr



*Corresponding author

§ Authors to whom inquiries concerning the X-ray structure should be directed.


NMR spectra of compounds **7**, **8**, **10**, **11**, **14**, **17**, **27** and **28**

Content


¹ H NMR spectrum of 7	S2
¹³ C NMR spectrum of 7	S3
¹ H NMR spectrum of 8	S4
¹³ C NMR spectrum of 8	S5
¹ H NMR spectrum of 10	S6
¹³ C NMR spectrum of 10	S7
¹ H NMR spectrum of 11	S8
¹³ C NMR spectrum of 11	S9
¹ H NMR spectrum of 14	S10
¹³ C NMR spectrum of 14	S11
¹ H NMR spectrum of 17	S12
¹³ C NMR spectrum of 17	S13
¹ H NMR spectrum of 27	S14
¹³ C NMR spectrum of 27	S15
¹ H NMR spectrum of 28	S16
¹³ C NMR spectrum of 28	S17





S2





7





S3







S4




S5










S8




11





S9





S10





S11








S13





S14



S15



S16



S17