



Synthesis of bi- and bis-1,2,3-triazoles by copper-catalyzed Huisgen cycloaddition: A family of valuable products by click chemistry

Zhan-Jiang Zheng^{*1}, Ding Wang¹, Zheng Xu¹ and Li-Wen Xu^{*1,2}

Review

Open Access

Address:

¹Key Laboratory of Organosilicon Chemistry and Material Technology of Ministry of Education, and College of Material, Chemistry and Chemical Engineering, Hangzhou Normal University, Hangzhou 310012, P. R. China and ²State Key Laboratory for Oxo Synthesis and Selective Oxidation, Lanzhou Institute of Chemical Physics, Chinese Academy of Sciences, Lanzhou, P. R. China

Email:

Zhan-Jiang Zheng^{*} - zzjiang78@hotmail.com; Zheng Xu - Zhengxu@hznu.edu.cn; Li-Wen Xu^{*} - liwenxu@hznu.edu.cn

* Corresponding author

Keywords:

bistriazoles; click chemistry; cycloaddition; homogeneous catalysis; oxidative coupling

Beilstein J. Org. Chem. **2015**, *11*, 2557–2576.

doi:10.3762/bjoc.11.276

Received: 31 August 2015

Accepted: 20 November 2015

Published: 11 December 2015

This article is part of the Thematic Series "Copper catalysis in organic synthesis".

Guest Editor: S. R. Chemler

© 2015 Zheng et al; licensee Beilstein-Institut.

License and terms: see end of document.


Abstract

The Cu(I)-catalyzed azide-alkyne cycloaddition reaction, also known as click chemistry, has become a useful tool for the facile formation of 1,2,3-triazoles. Specifically, the utility of this reaction has been demonstrated by the synthesis of structurally diverse bi- and bis-1,2,3-triazoles. The present review focuses on the synthesis of such bi- and bistriazoles and the importance of using copper-promoted click chemistry (CuAAC) for such transformations. In addition, the application of bitriazoles and the related CuAAAC reaction in different fields, including medicinal chemistry, coordination chemistry, biochemistry, and supramolecular chemistry, have been highlighted.

Introduction

Since its discovery by Huisgen and co-workers fifty years ago [1–4], the Huisgen cycloaddition of azides to alkynes has gained much attention due to its potential to yield a wide variety of triazoles with structurally diverse and functionalized groups, especially with respect to biological activity [5–8]. Originally, this transformation was typically carried out at high temperature and resulted in a mixture of the 1,4 and 1,5 regioisomers

(Scheme 1). Fortunately, representing a milestone in this field, the application of Cu(I) as the catalyst was reported by Sharpless and Meldal in 2002 [9,10]. In this work, the Huisgen reaction worked well under mild conditions, giving the desired triazoles with high yield, good regioselectivity, and quite high functional-group tolerances (Scheme 1). Since then, the so-called field of “click chemistry” has been extensively



Scheme 1: The synthesis of triazoles through the Huisgen cycloaddition of azides to alkynes.

investigated and recognized as an epoch-making progress in organic synthesis and green chemistry [11–15].

After many years of research, it was proven that the Cu(I)-catalyzed azide-alkyne cycloaddition (CuAAC reaction) could be performed under various conditions according to the need of click chemistry using the catalysis of various copper salts that generate Cu(I) sources *in situ* [16]. As a brief summary, the copper(I)-promoted click chemistry has the following features: (1) The most preferred methods for the formation of Cu(I) involve the use of CuSO₄ and a reducing agent in an aqueous solution. Most commonly, a large excess of sodium ascorbate with respect to the copper catalyst is the favored reducing agent, and a water/alcohol mixture is the favored solvent. This catalyst system combines the advantage of not requiring inert gas to prevent the Cu(I) from oxidation to Cu(II) during the reaction, and not requiring the tedious work-up and purification steps. (2) Other frequently used Cu(I) sources are CuI or CuBr. In general, CuI has been used in organic solvents such as CH₃CN, THF or toluene, and the active Cu(I) species was further stabilized by the addition of excess base. In addition, many other Cu(I) salts are used in CuAAC reactions owing to improved solubility or increased rate as compared to the CuSO₄/sodium ascorbate or CuI catalytic system. (3) The third type of Cu(I) source is generated by the oxidation of Cu metal. The Cu(0) species (found in forms such as turnings, wire, powder or nanoparticles) in the presence or absence of Cu(II) in aqueous media also provides the key active Cu(I) in some CuAAC reactions.


As an important supplement to the classic Huisgen cycloaddition promoted by copper catalysis, this review will deal with the copper-catalyzed syntheses of bi- and bistriazoles or their analogues by click chemistry, such as those linked directly or by spacers. The synthetic approaches for the preparation of bi- and bistriazoles are discussed in detail and their application is discussed briefly in each section. Accordingly, the following three types of bi- and bistriazoles will be primarily presented: (1) The 4,4'-linked symmetric or unsymmetrical bitriazoles. (2) The 5,5'-linked symmetric bitriazoles. (3) The bistriazoles formed through spacers from the dialkyne or diazide substrates.

Review

The synthesis of 4,4'-bitriazoles


The 4,4'-bitriazoles can be realized from a double CuAAC reaction between various sources of 1,3-butadiynes with the substituted azides. In general, two different methods have been developed for the construction of the 4,4'-bitriazoles: (1) The one-pot double CuAAC reaction of 1,3-butadiyne with azides. (2) Two successive CuAAC reactions with different or same azides that require the deprotection of the second reactive site to liberate another alkyne moiety.

In 2007, Monkowius et al. reported that the 4,4'-bitriazoles **3** could be synthesized by the two-fold click reaction between 1,3-butadiyne and substituted organic azides [17], and the reaction was catalyzed by a catalytic amount of CuI in acetonitrile in the presence of 2,4-lutidine. The desired, symmetrical 4,4'-bitriazoles **3** were obtained in good yield (76–82%, Scheme 2). However, the substrate 1,3-butadiyne (**1**) is difficult to handle because of its high reactivity and instability.



Scheme 2: The synthesis of symmetrically substituted 4,4'-bitriazoles.

Later in 2009, Fiandanese et al. proposed that unsymmetrically substituted 4,4'-bi-1,2,3-triazoles can be prepared in an easy manner [18]: starting from 1-trimethylsilyl-1,3-butadiyne (**4**) the 4-(silylalkynyl)-1,2,3-triazoles **5** can be obtained after reaction with substituted azides catalyzed by Cu(OAc)₂·H₂O (0.2 equiv). After the *in situ* deprotection with TBAF, followed by the CuI-catalyzed click reaction with another azide in THF in the presence of 1.2 equiv of 1,1,4,7,7-pentamethyldiethylenetriamine, the unsymmetrically 4,4'-bitriazoles **6** are obtained in good yield (52–86%, Scheme 3).




Simpson et al. also developed this three-step procedure (CuAAC–deprotection–CuAAC) into a one-pot fashion with moderate overall yield (34–49%) [19]. Similar to Fiandanese's strategy, Aizpurua et al. developed another synthetic method [20]: Starting with the CuAAC reaction of propargyl alcohol (**7**) with different azides, followed by the sequential Swern oxidation and Ohira–Bestmann homologation provided the ethynyltriazole intermediate **9**, finally another CuAAC resulted in the formation of unsymmetrical 4,4'-bi(1,2,3-triazole)s **10** (Scheme 4).

The synthesis of 5,5'-bitriazoles

Originally, in the research of the CuAAC reaction, the 5,5'-bitriazoles were usually considered as an undesired side product or impurity in the Huisgen cycloaddition. In general, they are the oxidative coupling product of the triazole-copper species. The 5,5'-bitriazoles were usually formed as the major product by the facilitation of the reaction conditions or controlled by the starting substrate.


In 2007, Burgess and Angell successfully developed an oxidative coupling method for the preparation of 5,5'-bitriazole [21]. In this work, they were able to perform this reaction of azides and terminal alkynes with moderate to high yield by using a 1:1 mixture of MeCN/2 M aqueous Na₂CO₃ solution at 25 °C for 18 h in the presence of a catalytic amount of CuSO₄ (10 mol %) and one equivalent of Cu powder (Scheme 5). Obviously, this



Scheme 5: The synthesis of 5,5'-bitriazoles.

method suffered from the drawback that a stoichiometric amount of Cu powder is required to achieve the highest activity.



In 2010, Nandurdikar et al. linked the two (or four) molecules of NO donor prodrugs together through the triazole spacers [22], which has potential application as NO-releasing materials. They first prepared the benzylidene-protected 2,2-di(azido-methyl)propane-1,3-diol containing the conformational strain.



They also investigated the click reactions with various alkynes under different conditions (Scheme 6), and found that: (1) when catalyzed with CuSO_4 /Na ascorbate in $\text{THF}/\text{H}_2\text{O}$, the reaction provided the normal bistriazoles **13** with moderate to good yield (60–75%). (2) By using CuI and DIPEA in acetonitrile to perform the reaction, the sequential, CuAAC oxidative coupling gave the cyclic 5,5'-bitriazoles **14** as the major product (44–74%). They demonstrated that both the reaction conditions

and the conformational effect are beneficial for the formation of the 5,5'-bitriazoles.

Similarly, Urbano and co-workers performed the CuAAC reaction of 1,1'-diazidoferrocene with ethynyl [5]-helicenequinone [23], and found the open chain bistriazolylphenyl-helicenequinone **17** could be obtained in good yield when CuSO_4 /sodium ascorbate was used in $\text{THF}/\text{H}_2\text{O}$ (Scheme 7).



However, the cyclic 5,5'-bitriazole **18** was achieved as the major product when CuI/Et₃N was used in CH₃CN.

In 2011, Cuevas-Yáñez and co-workers demonstrated that temperature was the key factor for the formation of 5,5'-bitriazole [24]. In that work, it was found that the desired bitriazoles could be obtained in low to moderate yield when the CuAAC reaction was performed at low temperature (-35°C) and high NaOH concentration. Subsequently, Jeon and co-workers reported a synthetic method for the construction of 5,5'-bitriazoles by using CuI together with 2 equiv of DIPEA [25]. The authors further demonstrated that the use of a base was also an important factor for the appropriate bitriazole yield. However, Jeon's method worked well only when the alkynes contained propargylic ethers and acetylenic amides (Scheme 8). Generally speaking, most of the above-mentioned oxidative coupling-click dimerizations merely provided the desired 5,5'-bitriazole in low yield when the alkynes or azides were linked directly with a hindered group or an aromatic moiety.

In 2012, Xu and co-workers disclosed that both the secondary and primary amine-functionalized polysiloxanes were good ligands for the copper-catalyzed Huisgen reaction of organic azides and alkynes [26]. This applied especially when the reaction was catalyzed by CuCl and mediated by the secondary amine-functionalized polysiloxanes at 0°C in CH₂Cl₂, where the symmetrically 5,5'-coupled bitriazole was obtained as the major product (Scheme 9). All the alkynes directly linked with the aromatic moiety provided the desired bitriazoles in moderate to good yield. We believe that the key copper mono-triazole intermediate formed after the first CuAAC reaction was stabilized by the secondary amine-functionalized polysiloxane, which further led to the formation of the bitriazole **22**. The chiral D-glucopyranosyl azide and phenylacetylene reacted under CuI and the secondary amine-functionalized poly-


siloxane catalytic system, where the desired chiral bitriazole **23** was obtained with good diastereoselectivity (85:15). Subsequently, the authors used this method to construct the novel, cyclic 5,5'-bitriazole **24** from binaphthol [27] (Scheme 10). Notably, this type of compound showed high selectivity over the recognition of I⁻, possibly due to the formation of a charge-transfer complex between the I⁻ and the electron-deficient triazole ring.

Bistriazoles formed through spacers



Bistriazole synthesis with dialkyne spacers

In the past years, bistriazoles with dialkyne spacers have gained significant attention for their potential application in supramolecular chemistry, pharmaceutical chemistry, biological chemistry and organometallic chemistry. The construction of bistriazoles from dialkynes is now well-developed, and three main methods are reported: (1) the one-pot CuAAC reactions of the terminal dialkynes with two equiv of the organic azides, providing the corresponding bistriazoles, which is the most popular method for the synthesis of bistriazoles with dialkyne spacers. (2) The second protocol is the sequential CuAAC-deprotection-CuAAC reaction for the construction of the bistriazoles. In general, the trialkylsilyl group was used as a temporary masking group for one of two alkyne moieties. Thus, this method provides the possibility of one molecule bearing two alkynes reacted with two different organic azides. (3) The third method involves the utilization of the substrate bearing two alkyne moieties with different reactivity in the successive Huisgen cycloaddition reactions: Huisgen reaction of the activated alkyne with the first azide and the CuAAC reaction of the nonactivated alkyne with another azide, leading to the corresponding bistriazoles.

In 2006, Aucagne and Leigh reported the synthesis of the TMS-alkyne and terminal-alkyne bis-functionalized tripeptide [28].



Scheme 8: The synthesis of 1,2,3-triazoles and 5,5'-bitriazoles from acetylenic amide.


**Scheme 9:** The amine-functionalized polysiloxane-mediated divergent synthesis of trizoles and bitriazoles.**Scheme 10:** The cyclic BINOL-based 5,5'-bitriazoles.

Here they used the CuAAC reaction of the terminal-alkyne-containing tripeptide **25** with the azide-containing dipeptide **26** in *t*-BuOH/H₂O, catalyzed by CuSO₄/sodium ascorbate, providing the first triazole-bearing intermediate (Scheme 11). They then performed the Ag(I)-catalyzed deprotection of the TMS-protected alkyne moiety, followed by another CuAAC reaction of the unmasked terminal alkyne with the second azide, giving the desired bistriazole-linked pseudononapeptides **28** in good yield (88–93%).


In 2009, Perumal and co-workers developed a one-pot methodology [29] for the multicomponent cycloaddition of sodium azide, benzyl bromides and various *N*-propargylated bis(indolyl)arylmethanes **30** catalyzed by CuI. It was proved that this multicomponent CuAAC reaction proceeded well in

PEG-400 among various solvents, mainly due to the hydrophilic and hydrophobic character of the PEG-400. As shown in Scheme 12, the CuAAC gave the desired bistriazoles **31** in good to excellent yield when electron-withdrawing groups were present in the benzyl bromides. Notably, the authors determined that all the compounds obtained showed potential biological activity.


Later, in 2011, Girard et al. demonstrated that the dialkynes with functional groups could react smoothly with various organic azides without the protection–deprotection step [30]. Initially, they first chose *N*-propargylpropiolamide **32** as the substrate and found the alkyne group with neighboring electron-withdrawing amide carbonyl was reacting exclusively with the organic azide under catalyst-free reaction conditions (with or without a solvent at room temperature) to give the mono-triazole intermediate **33** in good yield. Then the nonactivated terminal alkyne reacts with another azide, catalyzed by an Amberlyst A-21/CuI system in CH₂Cl₂ to form the second triazole ring with high yield (Scheme 13). In this work, the authors further extended this method to the propiolamide, which was derived from *meta*- and *para*-ethynylaniline, where both of the substrates worked well and the desired bistriazoles **34** could be obtained by a simple trituration and filtration procedure in good yield.



Scheme 11: The one-pot click–click reactions for the synthesis of bistriazoles.




Scheme 12: The synthesis of bis(indolyl)methane-derivatized 1,2,3-bistriazoles.




Scheme 13: The sequential, chemoselective preparation of bistriazoles.

The strain-promoted azide-alkyne cycloaddition (SPAAC) reaction could be well-performed without a Cu(I) catalyst. Such reactions are ideal for bioconjugation where no additional metal is required [31]. In 2012, Beal and co-workers incorporated the terminal alkyne and the activated cyclooctyne with two amide formation reactions [32], and then the SPAAC reaction was performed between the cyclooctyne **35** and the azides under mild conditions without the Cu(I). Notably, the CuAAC reaction between the terminal alkyne of **36** and various azides provided the desired bistriazoles **37** in good yield (Scheme 14).

Bistriazoles have potential application in the synthesis of surfactants after introducing the carbohydrate moiety. In 2012, Mohammed and co-workers selected the commercially available D-mannitol **38** as the starting material [33] in this reaction. After protection and introduction of the two alkyne groups to give dialkyne **39**, the copper-catalyzed cycloaddition occurred smoothly with the different long chain alkyl azides under the optimized reaction conditions. This provided the desired bistriazoles **40** in good yield (Scheme 15). It should be noted that the deprotection of bistriazoles **40** gives the gemini surfactants **41**



Scheme 14: The sequential SPAAC and CuAAC reaction for the preparation of bistriazoles.





Scheme 15: The synthesis of D-mannitol-based bistriazoles.

(with two hydrophobic arms and two hydrophilic heads) in high yield.

In 2012, Kaushik and co-workers prepared a series of ester-linked dialkynes from the acid dichloride and propargyl alcohol in the presence of DMAP [34]. Here, the CuAAC reaction between the dialkynes **42** and the azides provided the desired bistriazoles **43** in good yield (70–92%, Scheme 16). Interestingly, the antimicrobial activity studies revealed that compound **44** showed the highest activity against *B. subtilis* and *E. coli* due to the presence of a rigid pyridine nucleus. The authors further prepared various amide-linked bistriazoles by a three-component one-pot reaction of the amide-linked dialkynes, benzyl bromides and sodium azide catalyzed by $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ and sodium


ascorbate in DMF [35,36]. All the obtained compounds were evaluated for in vitro cytotoxicity against a panel of five human cancer cell lines, where compounds **45** and **46** displayed the highest and broadest spectrum activity against all five cancer cell lines under study (Scheme 16).

The formation of triazole functional groups can work as a powerful auxochrome. In 2010, Bunz and co-workers used the TMS-protected diethynylbenzothiadiazoles **48** and **50** as the source of dialkynes. The CuAAC reaction was carried out with ethylene-glycol-functionalized azide **47** in the presence of CuSO_4 and sodium ascorbate, providing the benzo-thiadiazole-based bistriazole (**49** and **51**, Scheme 17) [37]. With the aid of the hydrophilic character of the ethylene glycol group, both of



the bistriazoles can be endowed with water solubility, and can effectively bind Cu(II) and Ni(II) in water.

The thiocalix[4]arenes are sulfur-bridged analogs of calix[4]arenes, which have potential application in the molecular recognition of cationic, anionic or neutral molecules. In this regard, Yamato et al. incorporated two urea moieties possessing various aryl groups and two pyrene-appended triazole rings at the opposite side of the thiocalix[4]arene cavity [38]. The authors found that receptor **52** (Scheme 18) could selectively bind Cl⁻ through hydrogen bonding interaction with the urea NH protons, and **52** can also bind with Ag⁺ through complexation with the pyrene-appended bistriazole.



Scheme 18: The pyrene-appended thiocalix[4]arene-based bistriazole.

In homogeneous catalysis, functional ligands often play a key role in transition metal catalysis. Accordingly, the bistriazole derivatives could provide promising alternatives to bipyridine ligands because of their powerful nitrogen-centered coordination. In this context, Hao and co-workers have reported bistria-


zole-based N4 tetradentate ligands that were prepared by two CuAAC reactions in a one-pot procedure [39]. As shown in Scheme 19, these ligands exhibited good coordination properties to various metals, and the corresponding Mn(II) complexes showed good catalytic activity for the epoxidation of various aliphatic terminal olefins.

Very recently, Ulven and co-workers reported the synthesis of triazole-linked phenanthroline ligands. They were obtained by the following steps: (1) 1,10-phenanthroline-2,9-dicarbaldehyde (**56**) was treated with the Ohira–Bestmann reagent to provide the corresponding dialkyne **57**; (2) Dialkyne **57** was reacted with different azides catalyzed by the Cu(II)–TBTA complex and sodium ascorbate in a bi-phasic system of CH₂Cl₂/H₂O as the solvent, giving the desired bistriazoles **59** in good yield; (3) Deprotection of the N-Boc group with TFA, and the obtained primary amines was transformed to the corresponding primary guanidine or diisopropylguanidine analogs (**60** and **61**, Scheme 20) [40], which could be used as potential G4 DNA ligands with high selectivity over duplexed DNA.


Similar to the above strategies or methods, a number of researchers have developed various dialkyne substrates with varied spacers. As shown in Table 1, the reaction conditions are summarized for comparison, including the catalysts, the solvents, and the application of the bistriazoles.

Bistriazole synthesis with diazide spacers

The spacer-linked bistriazoles could also be prepared by the CuAAC reaction of the in situ generated diazides with substituted alkynes. There are mainly two types of methods for the construction of bistriazoles from diazides: (1) Starting from the substrate-containing good-leaving groups, the diazides could be generated in situ by nucleophilic addition with NaN₃, and then the double CuAAC reactions could give the desired bistriazoles. (2) Starting from a substrate containing one azide functionality and another suitable functional group, the CuAAC reaction was performed with a terminal alkyne to provide the monotriazole compound. Subsequently, the suitable functional group was transformed into the corresponding azide for the next CuAAC reaction to give the desired bistriazoles.



Scheme 19: The synthesis of triazole-based tetradentate ligands.

**Scheme 20:** The synthesis of phenanthroline-2,9-bistriazoles.**Table 1:** Previous reports on the copper-catalyzed Huisgen cycloaddition to bistriazoles with spacers.

| Cu source | Solvent | Spacer | Azide | Application |
|--|-----------------|--------|---|--------------|
| $CuSO_4 \cdot 5H_2O$, sodium ascorbate [41] | DMSO/ H_2O | | BnN_3 | – |
| bioClick conditions [42] | – | | $N_3-CH_2-CH_2-R$ R = OH or Me | Complexation |
| $CuSO_4 \cdot 5H_2O$, sodium ascorbate [43] | CH_2Cl_2/H_2O | | 6 azidosugars | – |
| A-21-CuI [44] | CH_2Cl_2 | | $R-N_3$, R = $Bn, CH_2CO_2Et, (CH_2)_3OAc, (CH_2)_3OH$ | – |

Table 1: Previous reports on the copper-catalyzed Huisgen cycloaddition to bistriazoles with spacers. (continued)

| | | | | |
|--|------------------------------------|--|---|--------------------------|
| $\text{Cp}^*\text{RuCl}(\text{COD})$ [45] | THF | | | Chemical sensors |
| CuSO_4 , sodium ascorbate [46] | $\text{H}_2\text{O}/t\text{-BuOH}$ | | 3 azides | Biological imaging |
| $[\text{Cu}(\text{CH}_3\text{CN})_4](\text{PF}_6)_2$ [47] | CH_2Cl_2 | | | Anion recognition |
| CuI [48] | THF | | | Receptors for fullerenes |
| $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, sodium ascorbate [49] | $\text{H}_2\text{O}/t\text{-BuOH}$ | | | Chemical sensors |
| $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, sodium ascorbate [50] | $\text{H}_2\text{O}/t\text{-BuOH}$ | | $\text{PhCH}_2\text{N}_3,$ $\text{F}_{13}\text{C}_6\text{N}_3$ | |
| CuSO_4 , sodium ascorbate [51] | $\text{H}_2\text{O}/t\text{-BuOH}$ | | | Chemical sensor |

Table 1: Previous reports on the copper-catalyzed Huisgen cycloaddition to bistriazoles with spacers. (continued)

















| | | | | |
|--|---|--|--|-------------------------|
| CuI [52] | – |  X = CH or N R = H or Me |  | Biological activity |
| CuSO ₄ ·5H ₂ O, sodium ascorbate [53] | CH ₂ Cl ₂ /H ₂ O |  | PhCH ₂ N ₃ | – |
| CuSO ₄ ·5H ₂ O, sodium ascorbate [54] | DMF/H ₂ O |  |  20 azides | Hemolytic activity |
| CuSO ₄ ·5H ₂ O, sodium ascorbate [55] | DMF/H ₂ O |  |  | Complexation |
| CuSO ₄ ·5H ₂ O, sodium ascorbate [56] | EtOH/H ₂ O |  |  n = 1, 2 | Antitubercular activity |
| CuSO ₄ ·5H ₂ O/Cu [57] | CH ₃ CN |  | PhCH ₂ N ₃  | – |
| CuSO ₄ , sodium ascorbate [58] | H ₂ O/t-BuOH |  |  | Biological activity |
| CuI [59] | THF/H ₂ O |  |  | Chemical sensor |

Table 1: Previous reports on the copper-catalyzed Huisgen cycloaddition to bistriazoles with spacers. (continued)


In 2007, Wang and co-workers demonstrated that the one-pot three-component reaction of *ortho*- and *meta*-bis(chloromethyl)benzene (**62**), sodium azide, and terminal alkynes, catalyzed by CuX in water could provide the corresponding 1,4-disubstituted bistriazoles **63** in excellent yield [64] (Scheme 21). They found that the bistriazole could be formed during the Huisgen reaction, in which the reaction was efficiently promoted by the catalytic amount of the Cu(I) salts. For

example, the three-component reaction could reach completion in the presence of only 0.2 mol % of Cu(I).

In 2010, Shreeve et al. reported that the diazides **65** could be generated in situ by the nucleophilic substitution of SF₅ and OTs groups with 3 equiv of NaN₃, followed by the CuAAC reaction with aliphatic alkynes, providing the desired bistriazoles **66** in moderate yield (Scheme 22) [65]. However, when



Scheme 21: The three-component reaction for the synthesis of bistriazoles.


**Scheme 22:** The one-pot synthesis of bistriazoles.


they choose the substrate SF_5 -ethylbromide, only trace amounts of the triazole-containing compound was obtained even after 18 h at 60 °C due to the poor leaving ability of the Br^- compared to the OTs group.

In 2011, Bundle et al. reported the double-click method for the formation of disymmetric bistriazoles [66]. The authors introduced the azide group (**69**) by the coupling of the protected amine functionality to the polymer substrate for the first CuAAC reaction (Scheme 23). Then, the treatment of the amine-containing mono-triazole intermediate **71** with the diazo transfer agent (imidazole-1-sulfonyl azide) was performed to convert the amine group into the corres-

ponding azide group, which provided a polymeric substrate for the second CuAAC reaction to give the desired bistriazoles (Scheme 23).

In 2009, Zhu and co-workers found that copper(II) acetate (Cu(OAc)_2) could catalyze the Huisgen alkyne-azide cycloaddition reactions without the addition of the reducing agents and could be produced in high yield when the substrate contains the chelating azide group [67]. Then they synthesized the bifunctional compounds with chelating azide groups and nonchelating azide groups (compounds **75**–**78**, Scheme 24) [68], by adding the Cu(OAc)_2 . This promoted the Huisgen cycloaddition of the chelating azide with the terminal alkyne, providing the mono-

**Scheme 23:** The synthesis of polymer-bearing 1,2,3-bistriazoles.

**Scheme 24:** The synthesis of bistriazoles via a sequential one-pot reaction.

triazole intermediates. Then, the introduction of the second alkyne, together with sodium ascorbate as the reducing agent, gave the desired bistriazoles in high yield. By mixing the two alkynes with different reactivities with the diazide, they developed a sequential one-pot method for the construction of the bistriazoles (**79**, **80**, etc.).

Notably, although there are many examples of the formation of spacer-linked bistriazoles from the diazide substrates, these bistriazole products were obtained with almost the same strategy as previously reported. These works are listed in Table 2 together with the reaction conditions and the applications of the corresponding bistriazoles.







Table 2: Previous reports on the copper-catalyzed Huisgen cycloaddition to bistriazoles with spacers.

| Cu source | Solvent | Spacer | Alkyne | Application |
|--|----------------------------------|--------|--------|-----------------|
| Cu(OAc) ₂ , sodium ascorbate [69] | H ₂ O/ <i>t</i> -BuOH | | | Complexation |
| CuI [70,71] | THF/H ₂ O | | | Chemical sensor |

Table 2: Previous reports on the copper-catalyzed Huisgen cycloaddition to bistriazoles with spacers. (continued)

| | | | | |
|--|-------------------------|--|--|------------------------------------|
| CuSO ₄ ·5H ₂ O, sodium ascorbate [72] | DMF/H ₂ O | | | Supermolecular chemistry |
| CuSO ₄ ·5H ₂ O, sodium ascorbate [73] | DMF | | | Cytotoxic activity |
| [74] | CHCl ₃ | | | Polymer chemistry |
| CuSO ₄ ·5H ₂ O, sodium ascorbate [75] | THF/H ₂ O | | | – |
| CuSO ₄ ·5H ₂ O, sodium ascorbate [76] | H ₂ O/t-BuOH | | | DNA binding |
| CuSO ₄ ·5H ₂ O, sodium ascorbate [77] | THF/H ₂ O | | | Receptor |
| CuSO ₄ ·5H ₂ O, sodium ascorbate [78] | DMF/H ₂ O | | | Molecular recognition |
| Cu(OAc) ₂ /Cu [79] | DMSO | | | Fluorescence brightening agents |
| $R = \text{CH}_3 \text{ or } \text{CH}_2\text{CH}(\text{CH}_3)_2$ | | | | |

Table 2: Previous reports on the copper-catalyzed Huisgen cycloaddition to bistriazoles with spacers. (continued)

| | | | | |
|---|--------------------------------------|---|--|------------------------|
| CuSO_4/Cu [80] | EtOH |  |  | Chemical sensor |
| CuI [81] | $\text{CH}_2\text{Cl}_2/\text{MeOH}$ |  $n = 1, 2$ |  | Electroactive receptor |
| Cu(OAc)_2 sodium ascorbate [82] | $t\text{-BuOH}$ |  |  | Biological activity |

Conclusion

During the past 15 years, the CuAAC reaction has become a powerful tool for the synthesis of a large number of 1,4-disubstituted 1,2,3-triazoles and has led to applications in almost every field of chemistry and biochemistry. In this review, we summarized the recent progress of the CuAAC reaction, together with various popular related reactions, which have unexpected potential to yield the 4,4'-, 5,5'-bitriazoles or spacer-linked bitriazoles. Nevertheless, as compared to the classic spacer-linked bitriazoles, the chemistry of 4,4'-, and 5,5'-bitriazoles (both their preparation and application) are still in their infancy. Although we restricted this review to describe only those having a bitriazole backbone, one can easily obtain a glimpse into the huge potential of bitriazoles in the broad sense when added to all other substitution possibilities. This topic, although already widely studied in the past years, is still continuously evolving and regularly brings new possibilities in click chemistry. We feel that this compilation will be beneficial to design practical approaches and better routes to improve the existing routes for the synthesis of synthetically useful bi- and

bistriazoles, and we expect that the pace of discovery of the application of bi- and bistriazoles in many fields will continue to increase for some time.

Acknowledgements

This project support by the National Natural Science Founder of China (No. 21173064, 51303043, and 21472031) and Zhejiang Provincial Natural Science Foundation of China (LR14B030001) is appreciated. Z. J. Zheng also thanks the financial support of the public project of Zhejiang Province (2014C31131).

References

- Huisgen, R. *Pure Appl. Chem.* **1989**, *61*, 613–628. doi:10.1351/pac198961040613
- Huisgen, R.; Szeimies, G.; Möbius, L. *Chem. Ber.* **1967**, *100*, 2494–2507. doi:10.1002/cber.19671000806
- Bastide, J.; Hamelin, J.; Texier, F.; Ven, V. Q. *Bull. Soc. Chim. Fr.* **1973**, 2555–2579.
- Bastide, J.; Hamelin, J.; Texier, F.; Ven, V. Q. *Bull. Soc. Chim. Fr.* **1973**, 2871–2887.

5. Alvarez, R.; Velazquez, S.; San-Felix, A.; Aquaro, S.; De Clercq, E.; Perno, C.-F.; Karlsson, A.; Balzarini, J.; Camarasa, M. J. *J. Med. Chem.* **1994**, *37*, 4185–4194. doi:10.1021/jm00050a015
6. Genin, M. J.; Allwine, D. A.; Anderson, D. J.; Barbachyn, M. R.; Emmert, D. E.; Garmon, S. A.; Graber, D. R.; Grega, K. C.; Hester, J. B.; Hutchinson, D. K.; Morris, J.; Reischer, R. J.; Ford, C. W.; Zurenko, G. E.; Hamel, J. C.; Schaadt, R. D.; Stapert, D.; Yagi, B. H. *J. Med. Chem.* **2000**, *43*, 953–970. doi:10.1021/jm990373e
7. Brockunier, L. L.; Parmee, E. R.; Ok, H. O.; Candelore, M. R.; Cascieri, M. A.; Colwell, L. F., Jr.; Deng, L.; Feeney, W. P.; Forrest, M. J.; Horn, G. J.; MacIntyre, D. E.; Tota, L.; Wyvatt, M. J.; Fisher, M. H.; Weber, A. E. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 2111–2114. doi:10.1016/S0960-894X(00)00422-4
8. Fan, W.-Q.; Katritzky, A. R. In *Comprehensive heterocyclic chemistry II*; Katritzky, A. R.; Rees, C. W.; Scriven, E. F. V., Eds.; Elsevier Science: Oxford, 1996; Vol. 4, pp 1–126.
9. Tornøe, C. W.; Christensen, C.; Meldal, M. J. *Org. Chem.* **2002**, *67*, 3057–3064. doi:10.1021/jo011148j
10. Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 2596–2599. doi:10.1002/1521-3773(20020715)41:14<2596::AID-ANIE2596>3.0.CO;2-4
11. Moses, J. E.; Moorhouse, A. D. *Chem. Soc. Rev.* **2007**, *36*, 1249–1262. doi:10.1039/b613014n
12. Finn, M. G.; Fokin, V. V. *Chem. Soc. Rev.* **2010**, *39*, 1231–1232. doi:10.1039/c003740k
13. Ganesh, V.; Sudhir, V. S.; Kundu, T.; Chandrasekaran, S. *Chem. – Asian J.* **2011**, *6*, 2670–2694. doi:10.1002/asia.201100408
14. Thirumurugan, P.; Matosiuk, D.; Jozwiak, K. *Chem. Rev.* **2013**, *113*, 4905–4979. doi:10.1021/cr200409f
15. Tang, W.; Becker, M. L. *Chem. Soc. Rev.* **2014**, *43*, 7013–7039. doi:10.1039/C4CS00139G
16. Meldal, M.; Tornøe, C. W. *Chem. Rev.* **2008**, *108*, 2952–3015. doi:10.1021/cr0783479
17. Monkowius, U.; Ritter, S.; König, B.; Zabel, M.; Yersin, H. *Eur. J. Inorg. Chem.* **2007**, 4597–4606. doi:10.1002/ejic.200700479
18. Fiandanese, V.; Bottalico, D.; Marchese, G.; Punzi, A.; Capuzzolo, F. *Tetrahedron* **2009**, *65*, 10573–10580. doi:10.1016/j.tet.2009.10.078
19. Doak, B. C.; Scanlon, M. J.; Simpson, J. S. *Org. Lett.* **2011**, *13*, 537–539. doi:10.1021/o102852z
20. Aizpurua, J. M.; Azcune, I.; Fratila, R. M.; Balentova, E.; Sagartzazu-Aizpurua, M.; Miranda, J. I. *Org. Lett.* **2010**, *12*, 1584–1587. doi:10.1021/o1003127
21. Angell, Y.; Burgess, K. *Angew. Chem., Int. Ed.* **2007**, *46*, 3649–3651. doi:10.1002/anie.200700399
22. Oladeinde, O. A.; Hong, S. Y.; Holland, R. J.; Maciag, A. E.; Keefer, L. K.; Saavedra, J. E.; Nandurdikar, R. S. *Org. Lett.* **2010**, *12*, 4256–4259. doi:10.1021/o101645k
23. del Hoyo, A. M.; Latorre, A.; Diaz, R.; Urbano, A.; Carreño, M. C. *Adv. Synth. Catal.* **2015**, *357*, 1154–1160. doi:10.1002/adsc.201400998
24. González, J.; Pérez, V. M.; Jiménez, D. O.; Lopez-Valdez, G.; Corana, D.; Cuevas-Yañez, E. *Tetrahedron Lett.* **2011**, *52*, 3514–3517. doi:10.1016/j.tetlet.2011.05.002
25. Kwon, M.; Jang, Y.; Yoon, S.; Yang, D.; Jeon, H. B. *Tetrahedron Lett.* **2012**, *53*, 1606–1609. doi:10.1016/j.tetlet.2012.01.069
26. Zheng, Z.-J.; Ye, F.; Zheng, L.-S.; Yang, K.-F.; Lai, G.-Q.; Xu, L.-W. *Chem. – Eur. J.* **2012**, *18*, 14094–14099. doi:10.1002/chem.201202472
27. Wang, C.-Y.; Zou, J.-F.; Zheng, Z.-J.; Huang, W.-S.; Li, L.; Xu, L.-W. *RSC Adv.* **2014**, *4*, 54256–54262. doi:10.1039/c4ra09589h
28. Augagne, V.; Leigh, D. A. *Org. Lett.* **2006**, *8*, 4505–4507. doi:10.1021/o1061657d
29. Damodiran, M.; Muralidharan, D.; Perumal, P. T. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 3611–3614. doi:10.1016/j.bmcl.2009.04.131
30. Elamari, H.; Meganem, F.; Herscovici, J.; Girard, C. *Tetrahedron Lett.* **2011**, *52*, 658–660. doi:10.1016/j.tetlet.2010.11.141
31. Kele, P.; Mező, G.; Achatz, D.; Wolfbeis, O. S. *Angew. Chem., Int. Ed.* **2009**, *48*, 344–347. doi:10.1002/anie.200804514
32. Beal, D. M.; Albrow, V. E.; Burslem, G.; Hitchen, L.; Fernandes, C.; Lapthorn, C.; Roberts, L. R.; Selby, M. D.; Jones, L. H. *Org. Biomol. Chem.* **2012**, *10*, 548–554. doi:10.1039/c1ob06398g
33. Mohammed, A. I.; Abboud, Z. H.; Alghanimi, A. H. O. *Tetrahedron Lett.* **2012**, *53*, 5081–5083. doi:10.1016/j.tetlet.2012.07.014
34. Lal, K.; Kumar, A.; Pavan, M. S.; Kaushik, C. P. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 4353–4357. doi:10.1016/j.bmcl.2012.05.008
35. Lal, K.; Kaushik, C. P.; Kumar, K.; Kumar, A.; Qazi, A. K.; Hamid, A.; Jaglan, S. *Med. Chem. Res.* **2014**, *23*, 4761–4770. doi:10.1007/s00044-014-1038-5
36. Lal, K.; Kaushik, C. P.; Kumar, A. *Med. Chem. Res.* **2015**, *24*, 3258–3271. doi:10.1007/s00044-015-1378-9
37. Brombosz, S. M.; Appleton, A. L.; Zappas, A. J., II; Bunz, U. H. F. *Chem. Commun.* **2010**, *46*, 1419–1421. doi:10.1039/b920484a
38. Tomiyasu, H.; Shigyo, N.; Ni, X.-L.; Zeng, X.; Redshaw, C.; Yamato, T. *Tetrahedron* **2014**, *70*, 7893–7899. doi:10.1016/j.tet.2014.08.068
39. Hao, E.; Wang, Z.; Jiao, L.; Wang, S. *Dalton Trans.* **2010**, *39*, 2660–2666. doi:10.1039/b922043g
40. Nielsen, M. C.; Larsen, A. F.; Abdiadir, F. H.; Ulven, T. *Eur. J. Med. Chem.* **2014**, *72*, 119–126. doi:10.1016/j.ejmchem.2013.11.027
41. Zhang, X.; Li, H.; You, L.; Tang, Y.; Hsung, R. P. *Adv. Synth. Catal.* **2006**, *348*, 2437–2442. doi:10.1002/adsc.200600404
42. Li, Y.; Huffman, J. C.; Flood, A. H. *Chem. Commun.* **2007**, 2692–2694. doi:10.1039/b703301j
43. Pereira, G. R.; Santos, L. J.; Luduvico, I.; Alves, R. B.; Pereira de Freitas, R. *Tetrahedron Lett.* **2010**, *51*, 1022–1025. doi:10.1016/j.tetlet.2009.12.050
44. Elamari, H.; Jlalia, I.; Louet, C.; Herscovici, J.; Meganem, F.; Girard, C. *Tetrahedron: Asymmetry* **2010**, *21*, 1179–1183. doi:10.1016/j.tetasy.2010.06.013
45. Ruan, Y.-B.; Yu, Y.; Li, C.; Bogliotti, N.; Tang, J.; Xie, J. *Tetrahedron* **2013**, *69*, 4603–4608. doi:10.1016/j.tet.2013.04.024
46. Zhou, M.; Zhang, X.; Bai, M.; Shen, D.; Xu, B.; Kao, J.; Ge, X.; Achilefu, S. *RSC Adv.* **2013**, *3*, 6756–6758. doi:10.1039/c3ra40441b
47. White, N. G.; Beer, P. D. *Org. Biomol. Chem.* **2013**, *11*, 1326–1333. doi:10.1039/c2ob27229f
48. Mulla, K.; Shaik, H.; Thompson, D. W.; Zhao, Y. *Org. Lett.* **2013**, *15*, 4532–4535. doi:10.1021/o1402093a
49. Midya, G. C.; Paladhi, S.; Bhowmik, S.; Saha, S.; Dash, J. *Org. Biomol. Chem.* **2013**, *11*, 3057–3063. doi:10.1039/c3ob40075a
50. Skarpas, H.; Osipov, S. N.; Vorob'eva, D. V.; Odintsev, I. L.; Lork, E.; Röschenthaler, G.-V. *Org. Biomol. Chem.* **2007**, *5*, 2361–2367. doi:10.1039/b705510b
51. Hemamalini, A.; Mohan Das, T. *New J. Chem.* **2013**, *37*, 2419–2425. doi:10.1039/c3nj00072a
52. Krim, J.; Taourirte, M.; Engels, J. W. *Molecules* **2012**, *17*, 179–190. doi:10.3390/molecules17010179
53. Singh, N.; Metla, B. P. R.; Elias, A. J. *J. Organomet. Chem.* **2012**, *717*, 99–107. doi:10.1016/j.jorganchem.2012.07.023

54. Singh, M. K.; Tilak, R.; Nath, G.; Awasthi, S. K.; Agarwal, A. *Eur. J. Med. Chem.* **2013**, *63*, 635–644. doi:10.1016/j.ejmech.2013.02.027
55. Scott, S. Ø.; Gavey, E. L.; Lind, S. J.; Gordon, K. C.; Crowley, J. D. *Dalton Trans.* **2011**, *40*, 12117–12124. doi:10.1039/c1dt0551e
56. Kumar, K.; Carrère-Kremer, S.; Kremer, L.; Guérardel, Y.; Biot, C.; Kumar, V. *Dalton Trans.* **2013**, *42*, 1492–1500. doi:10.1039/c2dt32148c
57. Huo, J.-P.; Lü, M.; Wang, Z.; Li, Y. *Chin. J. Chem.* **2012**, *30*, 2411–2422. doi:10.1002/cjoc.201200638
58. Jervis, P. J.; Moulis, M.; Jukes, J.-P.; Ghadbane, H.; Cox, L. R.; Cerundolo, V.; Besra, G. S. *Carbohydr. Res.* **2012**, *356*, 152–162. doi:10.1016/j.carres.2012.02.034
59. Xu, H.-R.; Li, K.; Liu, Q.; Wu, T.-M.; Wang, M.-Q.; Hou, J.-T.; Huang, Z.; Xie, Y.-M.; Yu, X.-Q. *Analyst* **2013**, *138*, 2329–2334. doi:10.1039/c3an36789d
60. Anthony, P.; Bashir, N.; Parveen, R. *Asian J. Pharm. Sci.* **2014**, *4*, 9–13. doi:10.15272/ajbps.v4i3.501
61. Asmus, S.; Beckendorf, S.; Zurro, M.; Mück-Lichtenfeld, C.; Fröhlich, R.; Mancheño, O. G. *Chem. – Asian J.* **2014**, *9*, 2178–2186. doi:10.1002/asia.201402237
62. Fischer, C.; Weber, E. *J. Inclusion Phenom. Macrocyclic Chem.* **2014**, *79*, 151–160. doi:10.1007/s10847-013-0338-6
63. Bryant, J. J.; Lindner, B. D.; Bunz, U. H. F. *J. Org. Chem.* **2013**, *78*, 1038–1044. doi:10.1021/jo3024889
64. Wang, Z.-X.; Zhao, Z.-G. *J. Heterocycl. Chem.* **2007**, *44*, 89–92. doi:10.1002/jhet.5570440115
65. Huang, Y.; Gard, G. L.; Shreeve, J. M. *Tetrahedron Lett.* **2010**, *51*, 6951–6954. doi:10.1016/j.tetlet.2010.10.149
66. Guiard, J.; Fiege, B.; Kitov, P. I.; Peters, T.; Bundle, D. R. *Chem. – Eur. J.* **2011**, *17*, 7438–7441. doi:10.1002/chem.201003414
67. Brotherton, W. S.; Michaels, H. A.; Simmons, J. T.; Clark, R. J.; Dalal, N. S.; Zhu, L. *Org. Lett.* **2009**, *11*, 4954–4957. doi:10.1021/o19021113
68. Yuan, Z.; Kuang, G.-C.; Clark, R. J.; Zhu, L. *Org. Lett.* **2012**, *14*, 2590–2593. doi:10.1021/o1300899n
69. Camp, C.; Dorbes, S.; Picard, C.; Benoit, E. *Tetrahedron Lett.* **2008**, *49*, 1979–1983. doi:10.1016/j.tetlet.2008.01.086
70. Hung, H.-C.; Cheng, C.-W.; Ho, I.-T.; Chung, W.-S. *Tetrahedron Lett.* **2009**, *50*, 302–305. doi:10.1016/j.tetlet.2008.10.147
71. Hung, H.-C.; Cheng, C.-W.; Wang, Y.-Y.; Chen, Y.-J.; Chung, W.-S. *Eur. J. Org. Chem.* **2009**, 6360–6366. doi:10.1002/ejoc.200900987
72. Wei, P.; Yan, X.; Li, J.; Ma, Y.; Yao, Y.; Huang, F. *Tetrahedron* **2012**, *68*, 9179–9185. doi:10.1016/j.tet.2012.09.018
73. Jurášek, M.; Džubák, P.; Sedláč, D.; Dvořáková, H.; Hajdúch, M.; Bartúněk, P.; Drašar, P. *Steroids* **2013**, *78*, 356–361. doi:10.1016/j.steroids.2012.11.016
74. Isaacman, M. J.; Corigliano, E. M.; Theogarajan, L. S. *Biomacromolecules* **2013**, *14*, 2996–3000. doi:10.1021/bm400940h
75. Rajesh, R.; Periyasami, G.; Raghunathan, R. *Tetrahedron Lett.* **2010**, *51*, 1896–1898. doi:10.1016/j.tetlet.2010.02.020
76. Kamal, A.; Shankaraiah, N.; Reddy, C. R.; Prabhakar, S.; Markandeya, N.; Srivastava, H. K.; Sastry, G. N. *Tetrahedron* **2010**, *66*, 5498–5506. doi:10.1016/j.tet.2010.05.003
77. Romero, T.; Orenes, R. A.; Tárraga, A.; Molina, P. *Organometallics* **2013**, *32*, 5740–5753. doi:10.1021/om4002457
78. Karuturi, R.; Al-Horani, R. A.; Mehta, S. C.; Gailani, D.; Desai, U. R. *J. Med. Chem.* **2013**, *56*, 2415–2428. doi:10.1021/jm301757v
79. Huo, J.-P.; Luo, J.-C.; Wu, W.; Xiong, J.-F.; Mo, G.-Z.; Wang, Z.-Y. *Ind. Eng. Chem. Res.* **2013**, *52*, 11850–11857. doi:10.1021/ie401149p
80. Huang, H.-J.; Fang, H.-Y.; Chir, J.-L.; Wu, A.-T. *Luminescence* **2011**, *26*, 518–522. doi:10.1002/bio.1263
81. Mames, I.; Wawrzyniak, U. E.; Woźny, M.; Bilewicz, R.; Korybut-Daszkiewicz, B. *Dalton Trans.* **2013**, *42*, 2382–2391. doi:10.1039/c2dt32386a
82. Westermann, B.; Dörner, S.; Brauch, S.; Schaks, A.; Heinke, R.; Stark, S.; van Delft, F. L.; van Berkel, S. S. *Carbohydr. Res.* **2013**, *371*, 61–67. doi:10.1016/j.carres.2013.02.003

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the *Beilstein Journal of Organic Chemistry* terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at:

[doi:10.3762/bjoc.11.276](http://doi.org/10.3762/bjoc.11.276)