



Supporting Information

for

Preparation and in vivo evaluation of glyco-gold nanoparticles carrying synthetic mycobacterial hexaarabinofuranoside

Gennady L. Burygin, Polina I. Abronina, Nikita M. Podvalnyy, Sergey A. Staroverov, Leonid O. Kononov and Lev A. Dykman

Beilstein J. Nanotechnol. **2020**, *11*, 480–493. doi:10.3762/bjnano.11.39

Results of ELISA of *M. bovis*, *M. phlei*, *M. smegmatis* and *E. coli* cell suspensions




Figure S1: Results of ELISA of *M. bovis*, *M. phlei*, *M. smegmatis* and *E. coli* cell suspensions (10^8 cells·mL⁻¹) with antisera against Ara₆C₂NH₂-GNPs **3** (A), against Ara₆C₂EG₇NH₂-GNPs **4** (B), against intact serum of the rabbit used for immunization (C) or antibodies against LPS of *Azospirillum brasilense* Sp7 generated in the presence of CFA (D) [1]. Error bars indicate the observed confidence intervals at $p < 0.05$.

References

1. Konnova, O. N.; Boiko, A. S.; Burygin, G. L.; Fedonenko, Y. P.; Matora, L. Y.; Konnova, S. A.; Ignatov, V. V. *Microbiology (London, U. K.)* **2008**, 77, 305–312.
doi:10.1134/S0026261708030090