



Synthesis and conformational analysis of pyran inter-halide analogues of D-talose

Olivier Lessard, Mathilde Grosset-Magagne, Paul A. Johnson and Denis Giguère*

Full Research Paper

Open Access

Address:

Département de Chimie, 1045 av. De la Médecine, Université Laval,
Québec City, Qc, G1V 0A6, PROTEO, Canada

Email:

Denis Giguère* - denis.giguere@chm.ulaval.ca

* Corresponding author

Keywords:

organofluorine; pyran inter-halide; solid-state conformation;
solution-state conformation

Beilstein J. Org. Chem. **2024**, *20*, 2442–2454.

<https://doi.org/10.3762/bjoc.20.208>

Received: 05 June 2024

Accepted: 11 September 2024

Published: 27 September 2024

This article is part of the thematic issue "Organofluorine chemistry VI".

Guest Editor: D. O'Hagan



© 2024 Lessard et al.; licensee Beilstein-Institut.
License and terms: see end of document.

Abstract

In this work, we describe the synthesis of halogenated pyran analogues of D-talose using a halo-divergent strategy from known 1,6-anhydro-2,3-dideoxy-2,3-difluoro-β-D-mannopyranose. In solution and in the solid-state, all analogues adopt standard ⁴C₁-like conformations despite 1,3-diaxial repulsion between the F2 and the C4 halogen. Moreover, the solid-state conformational analysis of halogenated pyrans reveals deviation in the intra-annular torsion angles arising from repulsion between the axial fluorine at C2 and the axial halogen at C4, which increases with the size of the halogen at C4 (F < Cl < Br < I). Crystal packing arrangements of pyran inter-halides show hydrogen bond acceptor and nonbonding interactions for the halogen at C4. Finally, density functional theory (DFT) calculations corroborate the preference of talose analogues to adopt a ⁴C₁-like conformation and a natural bonding orbital (NBO) analysis demonstrates the effects of hyperconjugation from C–F antibonding orbitals.

Introduction

Polyfluorinated pyran analogues of carbohydrates have attracted attention over the years. This class of glycomimetics has great biological potential with useful applications [1–7]. What about other halogens? Pyran inter-halide analogues of carbohydrates were rarely investigated as new tools in glycobiology [8]. This is surprising since the incorporation of halogens can improve cellular uptakes and enhance membrane binding and permeation [9–11]. In addition, halogen bonding is an impor-

tant interaction in biological systems [12–17] and the beneficial effect of the chloro substituent has been reviewed recently [18].

As a result, there is a lack of efficient synthetic strategies to access multivincinal inter-halide stereocenters (i.e., contiguous chiral halides: F, Cl, Br, I) [19]. Only a handful of natural product syntheses have been reported [20,21], despite the promising biological activity of these unique inter-halides [22]. For our

part, we recently reported the synthesis of contiguous inter-halide-bearing stereocenters using a Chiron approach from levoglucosan **1** (Figure 1a) [23]. Allopyranose inter-halides **4** incorporating the 2,3-*cis*, 3,4-*cis* relationship for the halogens were prepared via intermediates **2** and **3** from levoglucosan (**1**). Compounds **4** were the starting point to complex 2,3,4-trihalo-hexanetriols and 2,3,4,5-tetrahalohexanediols. Conformational analysis and lipophilicities of the latter compounds were determined and trihalogenated alkanes were incorporated into piperidines of pitolisant [23]. This work was an extension of our synthetic routes to multivincinal organofluorines to unveil some of their unique properties [24–30], such as the solution-state conformation of diastereomeric polyfluorohexitols [31].

Herein, we report the synthesis of pyran inter-halide analogues of D-talopyranose **6**, integrating also the 2,3-*cis*, 3,4-*cis* relationship for the halogens, from known intermediate **5** (Figure 1b) [24]. The solution and the solid-state conformational analysis were supplemented with DFT calculations to understand key conformational drivers. This study adds more data to the field of nuclear magnetic resonance (NMR) spectroscopy of polyhalogenated molecules. It should be noted that the NMR predictions of such compounds remain very challenging [32].

Results and Discussion

Our recent discovery that the nature of halogen atoms can have a large impact on conformation and lipophilicity motivated us to investigate novel pyran inter-halides [23]. We used a halo-divergent route starting from the known 1,6-anhydro-2,3-dideoxy-2,3-difluoro-β-D-mannopyranose (**5**) readily accessible from levoglucosan (**1**, Scheme 1) [24]. Activation of the C4 hydroxy group as triflate and direct treatment with a nucleophilic halogen furnished intermediates **8–11**. The latter com-

pounds proved to be difficult to purify, therefore we were compelled to proceed directly to the next step. Cleavage of the 1,6-anhydro bridges was achieved under acetolysis conditions providing halogenated talopyranoses **12–15** in good yield over 3 steps as α anomers.

Luckily, inter-halides **13–15** were crystalline, allowing the absolute configuration to be confirmed by single-crystal diffraction analysis (see below) [33]. Unfortunately, trifluorinated analogue **12** was not crystalline. Thus, we removed the acetyl protecting groups and generated the corresponding *p*-bromobenzoate derivative **17** to obtain suitable crystalline material [24,34].

In order to decipher the key physical properties of complex pyran inter-halides, we performed ^{19}F NMR analysis of halogenated talose analogues **12–15** (Figure 2). First, all analogues adopt standard $^4\text{C}_1$ -like conformations. Comparison of the vicinal and geminal coupling constants for each organohalogen suggests that there is little change in the conformations (although there is an increasing chair distortion for larger halogens, see below). Because F3 is adjacent to the C4 halogen, ^{19}F resonance of F3 occurs at lower field than F2 for analogues **13–15**. There is a strong increase in chemical shift of F3 depending on the incorporated halogen on the pyran core at C4: –208.33 ppm for **12** (fluorine), –197.95 ppm for **13** (chlorine), –192.80 ppm for **14** (bromine), and –184.56 ppm for **15** (iodine). Similarly, the increase in chemical shift of F2 is smaller as exemplified with an upfield shift of –205.46 ppm for **12** to –200.55 ppm for **15**. Talopyranose analogues **12–15** incorporate a 2,3-*cis*, 3,4-*cis* relationship for the halogens. We previously prepared a small set of trihalogenated allopyranose analogues that also included the 2,3-*cis*, 3,4-*cis* relationship for





Figure 1: Synthesis of trihalogenated pyrans: a) Chiron approach to multivincinal inter-halide derived from allopyranoses; b) synthesis and conformational analysis of pyran inter-halide analogues of D-talose integrating the 2,3-*cis*, 3,4-*cis* relationship for the halogens (this work).



Scheme 1: Synthesis of halogenated talopyranose analogues **13–15**, and **17** that include a 2,3-*cis*, 3,4-*cis* relationship for the halogens.




Figure 2: Direct comparison of ^{19}F resonances of halogenated talose analogues **12–15** (^{19}F NMR; 470 MHz, CDCl_3).

the halogens (Figure 1a) [23]. ^{19}F NMR analysis of halogenated allopyranose analogues can be found in Supporting Information File 1.

Our interest in the conformation of organohalogens motivated us to compare the solid-state conformation of halogenated pyrans **13**–**15**, and **17** [24] with α -D-talose **18** [35] (Figure 3). The crystallographic data and structural refinement details for the crystal structures can be found in Supporting Information File 1. As tosyl and benzoate groups are essential for the crystallinity of multivincinal organofluorines they influence the solid-state conformations [31,36–38]. Thus, information drawn from the crystallographic data of compound **17** might be influenced by benzoate groups. We included compound **17** in our comparative analysis in any case. All structures adopt a standard $^4\text{C}_1$ -like conformation in the solid-state. This conformation occurs despite 1,3-diaxial repulsion between the F2 and the C4 halogen. The 1,3-diaxial repulsion between 2 fluorine atoms

have been reported in recent years [24,39], however, the 1,3-diaxial repulsion between fluorine and other halogens is quite uncommon [40,41]. As for the C5–C6 rotamer, all analogues exhibit a *gt* conformation except for trifluorinated **17**, which possesses a *tg* conformation. An axial substituent at C4 generally leads to a *gt* conformation [42,43], with some exceptions [24]. Bond distances, bond angles, torsion angles, and key interatomic distances are listed in Tables 1–4. It is important to point out that these results compare well with previous analysis of polyfluorinated carbohydrates [44,45].

The C–C bond lengths within the pyran rings of halogenated analogues are between 1.50 and 1.54 Å, which is similar to native talose (**18**, 1.52–1.53 Å) (Table 1, entries 1–4). However, all specified bond lengths within the pyran rings are shorter for halogenated analogues compared to α -D-talose, except for the C3–C4 bond of compound **15**. This can be explained by the adjacent repulsion between CF₃ with the Cl₄ group. Next, it has




Table 1: Selected bond distances for compounds **13–15**, **17**, and **18**.

Entry	Bonds	Distances (Å)				
		Talose (18) ^a	17	13	14	15
1	C1–C2	1.5316	1.518(3)	1.526(2)	1.523(3)	1.53(1)
2	C2–C3	1.5234	1.506(3)	1.513(2)	1.508(3)	1.50(2)
3	C3–C4	1.5300	1.508(4)	1.522(2)	1.520(3)	1.54(2)
4	C4–C5	1.5325	1.524(3)	1.527(2)	1.523(3)	1.53(1)
5	C5–C6	1.5127	1.525(3)	1.507(2)	1.513(3)	1.50(1)
6	C5–O5	1.4489	1.432(3)	1.438(2)	1.435(2)	1.44(1)
7	O5–C1	1.4380	1.399(3)	1.403(2)	1.403(2)	1.40(1)
8	C1–O1	1.4028	1.442(3)	1.429(2)	1.431(2)	1.44(1)
9	O1–C(O)	na	1.361(3)	1.372(2)	1.374(3)	1.40(1)
10	C2–F2	1.4228 ^b	1.393(3)	1.402(2)	1.407(2)	1.39(1)
11	C3–F3	1.4212 ^c	1.397(3)	1.393(2)	1.396(2)	1.39(1)
12	C4–X4	1.4279 ^d	1.395(3)	1.797(2)	1.956(2)	2.11(1)

^aReference [35]; ^bC2–O2; ^cC3–O3; ^dC4–O4.

been reported that the C1–O1 bond lengths are shorter than the O5–C1 for α anomers [42,46,47]. Talopyranose (**18**) follows this trend, but not the halogenated analogues (Table 1, entries 7 and 8). Also, the exocyclic C1–O1 bond lengths of compounds **13–15** and **17** are in average 0.033 Å longer than native talose (**18**). As expected, all the C–F bond lengths are shorter than the corresponding C–OH bond lengths (Table 1, entries 10–12) [48]. The C2–F2 bond lengths are in average 0.025 Å shorter than the C2–OH bond and the C3–F3 bond lengths are in average 0.027 Å shorter than the C3–OH bond. Similarly, for compound **13–15**, the C4–X bond lengths are longer than the C4–OH bond of native talose (1.43 Å): C4–Cl: 1.80 Å, C4–Br: 1.96 Å, and C4–I: 2.11 Å.

Table 2 shows the selected bond angles for compounds **13–15**, **17**, and **18**. All the bond angles of halogenated analogues are larger by 0.1–4.07° than talose (**18**). As such, the H2–C2–F2 bond angles are similar for compounds **13–15** and **17** (109.69–109.78°), but significantly larger than the H2–C2–O2 bond angle of talose (105.87°). Moreover, the H3–C3–F3 bond angles slightly decrease according to the nature of the atom at C4 (F: 107.97°; Cl: 107.67°; Br: 107.47°; I: 107.46°), as compared with talose (108.85°). As for the H4–C4–X4 bond angles, the angles are similar for talose and the trifluorinated analogues: 109.33° and 109.36°, respectively. However, there is a bond angle narrowing for the other analogues (Cl: 108.07°; Br: 107.94°; I: 107.78°). Finally, the angles involving the *exo*-

Table 2: Selected bond angles for compounds **13–15**, **17**, and **18**.

Entry	Bonds	Angles (°)				
		Talose (18) ^a	17	13	14	15
1	C1–C2–C3	109.51	110.2(2)	111.9(1)	111.9(2)	111.9(9)
2	C2–C3–C4	110.43	113.7(2)	114.0(1)	114.5(2)	114.1(9)
3	C3–C4–C5	107.80	109.9(2)	108.3(1)	108.3(2)	107.9(8)
4	C4–C5–O5	109.92	111.8(2)	111.6(1)	112.2(1)	112.0(8)
5	C5–O5–C1	113.68	114.9(2)	114.7(1)	114.7(1)	113.9(8)
6	O5–C1–C2	110.29	113.0(2)	112.2(1)	112.3(2)	112.7(9)
7	O1–C1–O5	111.87	111.1(2)	111.3(1)	110.9(2)	111.4(8)
8	O1–C1–C2	107.98	105.5(2)	106.8(1)	106.9(2)	106.6(8)
9	C1–O1–C(O)	na	116.2(2)	115.4(1)	115.3(2)	115.2(8)
10	C1–C2–F2	109.78 ^b	106.9(2)	105.4(1)	105.2(1)	104.6(8)
11	C3–C2–F2	112.49 ^c	110.5(2)	110.4(1)	110.5(2)	110.9(9)

Table 2: Selected bond angles for compounds **13–15**, **17**, and **18**. (continued)

12	C2–C3–F3	107.54 ^d	109.3(2)	109.5(1)	109.6(1)	109.9(9)
13	C4–C3–F3	113.36 ^e	109.7(2)	110.1(1)	110.0(1)	110.1(8)
14	C3–C4–X4	108.18 ^f	109.7(2)	112.1(1)	112.1(1)	111.9(7)
15	C5–C4–X4	111.14 ^g	109.2(2)	112.1(1)	112.5(1)	113.5(7)
16	H2–C2–F2	105.87 ^h	109.75	109.69	109.72	109.78
17	H3–C3–F3	108.85 ⁱ	107.97	107.67	107.47	107.46
18	H4–C4–X4	109.33 ^j	109.36	108.07	107.94	107.78

^aReference [35]; ^bC1–C2–O2; ^cC3–C2–O2; ^dC2–C3–O3; ^eC4–C3–O3; ^fC3–C4–O4; ^gC5–C4–O4; ^hH2–C2–O2; ⁱH3–C3–O3; ^jH4–C4–O4.

anomeric oxygen (O1–C1–O5) are similar with a difference of about 1° between the larger (compound **17**) and smaller (compound **14**) angle.

As stated above, all analogues exhibit a *gt* conformation except for compound **17**, which is a *tg* conformer. This information could also be extracted from Table 3 by looking at the O5–C5–C6–O6 torsion angles (Table 3, entry 1). Table 3 also highlights that there are significant intra-annular torsion angles for halogenated analogues. There are reductions in the C1–C2–C3–C4 torsion angles for the halogenated pyrans as compared to compound **18** (–56.58°) (Table 3, entry 4). The decrease depends on the size of the halogen at C4 (F: –49.4°; Cl:

–46.9°; Br: –46.5°; I: –46°). There is also a reduction in the C2–C3–C4–C5 torsion angles of about 8.4° in average for compound **13–15** and **17** falling outside the range of an ideal pyran ring (Table 3, entry 5) [35]. However, the C4–C5–O5–C1 torsion angles are similar for all compounds except for compound **17** (Table 3, entry 7). The deviation in the intra-annular torsion angles clearly arise from repulsion of the axial fluorine at C2 and the axial halogen at C4 as exemplified with the H3–C3–C2–F2 and the X4–C4–C3–H3 torsion angles being smaller than the expected 180° (Table 3, entry 16 and 17). The 1,3-diaxial repulsion leads to a deviation from parallel alignment as shown in Table 4. The Newman projections of the halogenated analogues show deviations from parallel alignment for

Table 3: Selected torsion angles for compounds **13–15**, **17**, and **18**.

Entry	Bonds	Torsion angles (°)				
		Talose 18 ^a	17	13	14	15
1	O5–C5–C6–O6	70.35	–178.3(2)	75.4(1)	74.9(2)	75(1)
2	O5–C1–O1–C(O)	na	94.7(2)	89.0(2)	88.7(2)	88(1)
3	C2–C1–O1–C(O)	na	–142.5(2)	–148.1(1)	–148.5(2)	–148.4(9)
4	C1–C2–C3–C4	–56.58	–49.4(3)	–46.9(2)	–46.5(2)	–46(1)
5	C2–C3–C4–C5	57.88	50.1(3)	50.0(2)	49.0(2)	49(1)
6	C3–C4–C5–O5	–58.49	–51.6(3)	–55.0(2)	–53.9(2)	–55(1)
7	C4–C5–O5–C1	60.88	56.4(2)	60.5(2)	59.8(2)	61(1)
8	C5–O5–C1–C2	–58.73	–56.1(2)	–55.5(2)	–55.3(2)	–57(1)
9	O5–C1–C2–C3	55.31	51.0(3)	47.6(2)	47.6(2)	48(1)
10	O5–C1–C2–F2	–68.66 ^b	–69.0(2)	–72.4(1)	–72.4(2)	–72(1)
11	C1–C2–C3–F3	179.26 ^c	–172.3(2)	–170.7(1)	–170.7(1)	–170.6(8)
12	F2–C2–C3–C4	65.78 ^d	68.5(3)	70.2(2)	70.4(2)	70(1)
13	F3–C3–C4–C5	178.62 ^e	172.8(2)	173.5(1)	173.0(1)	173.1(8)
14	C2–C3–C4–X4	–62.40 ^f	–70.0(3)	–74.2(2)	–75.6(2)	–77(1)
15	X4–C4–C5–O5	59.91 ^g	68.8(2)	69.2(1)	70.5(2)	70.0(9)
16	X4–C4–C3–H3	179.05 ^h	170.2	166.4	165.0	164.4
17	H3–C3–C2–F2	–175.98 ⁱ	–171.7	–170.4	–170.3	–170.9

^aReference [35]; ^bO5–C1–C2–O2; ^cC1–C2–C3–O3; ^dO2–C2–C3–C4; ^eO3–C3–C4–C5; ^fC2–C3–C4–O4; ^gO4–C4–C5–O5; ^hO4–C4–C3–H3; ⁱH3–C3–C2–O2.

the C2–F and C4–X substituents of 12.08° for **17**, 17.08° for **13**, 18.18° for **14**, and 18.59° for **15** (talopyranose, **18**: 5.92°). This deviation is responsible for the distance between F2 and the halogen at C4. Table 5 highlights key interatomic distances for all analogues. As such, the F2...X4 distance increases with the size of the C4 halogen (F < Cl < Br < I): 2.82 Å for **17**, 3.06 Å for **13**, 3.14 Å for **14**, and 3.23 Å for **15**. Another interesting feature can be drawn from Table 5. As such, intramolecular F2...F3, F2...X4, and F3...X4 contacts are smaller than the sum of the Van der Waals radii [49]. Taking together, all these data clearly demonstrate that the nature of one halogen can have an impact on the solid-state conformation of halogenated pyrans.


We also evaluated the Cremer–Pople ring puckering parameters (Table 6) [50]. For pyranoid rings, these parameters take the form of a spherical polar coordinate set, Q , θ , and ϕ , which define the point P, on a sphere of radius Q [51]. The smaller puckering amplitude (Q) values for the halogenated analogues indicate a flattened ring in comparison to the non-halogenated

Table 6: Cremer–Pople ring puckering amplitudes (Q), theta (θ) and phi (ϕ) parameters.

	Talose (18)	17	13	14	15
Q (Å)	0.588	0.514	0.521	0.514	0.523
θ (°)	2.976	1.724	6.936	6.439	7.248
ϕ (°)	233.600	315.204	299.454	305.438	307.711

compound. The puckering amplitude for an ideal cyclohexane chair, with C–C bond lengths of 1.54 Å, is 0.63 Å [50]. The azimuthal angle (θ) represents the distortion of the ring. For pyranose rings, an azimuthal angle of $\theta = 0^\circ$ represents a perfect 4C_1 chair, and an angle of $\theta = 180^\circ$ is the 1C_4 chair. The distortion of the chair conformation increases with the size of the halogen at C4. Surprisingly, the trifluorinated analogue is less distorted than the non-halogenated talopyranose. The meridian angle (ϕ) indicates the nature of the distortion. The distortion of the trifluorinated analogue is in a direction somewhat between an E_5 conformation ($\phi \approx 300^\circ$) and a 0H_5 conformation ($\phi \approx$

Table 4: 1,3-Diaxial repulsion between C2–F and C4–X bonds for **13–15** and **17**.^a

Entry	Angles (°)				
		17 (X = F, R = <i>p</i> -BrBz)	13 (X = Cl, R = Ac)	14 (X = Br, R = Ac)	15 (X = I, R = Ac)
1	α_1	5.99	6.77	7.05	7.06
2	α_2	6.09	10.31	11.13	11.53
3	$\alpha_1 + \alpha_2$	12.08	17.08	18.18	18.59

^aFor α -talopyranose **18**: $\alpha_1 = 5.58^\circ$, $\alpha_2 = 0.34^\circ$, and $\alpha_1 + \alpha_2 = 5.92^\circ$ (reference [35]).

Table 5: Key interatomic distances (intramolecular) for **13–15**, **17**, and **18**.

Entry	Atoms	Distances (Å)				
		Talose (18) ^a	17	13	14	15
1	O1...F2	3.6141 ^b	3.560(2)	3.552(2)	3.557(2)	3.55(1)
2	O1...F3	4.1700 ^c	4.194(2)	4.288(2)	4.285(2)	4.28(1)
3	F2...F3	2.8154 ^d	2.732(3)	2.735(2)	2.741(2)	2.746(9)
4	F2...X4	2.6546 ^e	2.817(2)	3.056(1)	3.143(1)	3.228(6)
5	F3...X4	2.8517 ^f	2.714(2)	2.968(1)	3.052(1)	3.160(6)

^aReference [35]; ^bO1...O2; ^cO1...O3; ^dO2...O3; ^eO2...O4; ^fO3...O4.

330°). The other trihalogenated analogues are distorted towards an E_5 conformation ($\varphi \approx 300^\circ$). The distortion of the non-halogenated talopyranose is in the direction on an 4E conformation ($\varphi \approx 240^\circ$).

The solid-state conformation of each of the pyran inter-halides **13–15** is unique. One would expect that compounds **13–15** would have distinct crystal packing arrangements based on the nature of the halogen. On the contrary, all analogues adopt a similar stacking pattern. Figure 4 shows the packing arrangement for compound **15** and the crystal packing of compound **13** and **14** can be found in the Supporting Information File 1. The halogens are on the same side of the pyran ring, thus increasing the overall molecular dipole moment (see Supporting Information File 1). This allows intermolecular C–X...H–C interactions responsible, in part, for the solid-state ordering [52]. Individual pyrans stack on top of one another in a manner consistent with electrostatic interactions with halogens facing H3, H4, and H5 (Figure 4a). As such, some intermolecular H...X bond distances and angles for compound **13–15** are listed in Table 7. Solid-state intermolecular interactions involving fluorine atoms have been well documented over the years for carbohydrate analogues [24,44,53] or other organofluorines [54]. In our case, there is a number of C–F...H–C interactions with F2 and H4 (**13**: $d = 2.271 \text{ \AA}$, **14**: $d = 2.270 \text{ \AA}$, and **15**: $d = 2.356 \text{ \AA}$) and with F3 and H4 (**13**: $d = 2.867 \text{ \AA}$, **14**: $d = 2.849 \text{ \AA}$, and **15**: $d = 2.886 \text{ \AA}$).

Does the halogen at C4 contribute to the stabilization within the crystal lattice? To answer this question, we have to look at the behavior of halogens as hydrogen bond acceptors ($X\cdots H$) and nonbonding interactions ($X\cdots O/N/S$). For C–X, a σ -hole arises when a valence electron is pulled into the σ -molecular orbital resulting in an electropositive crown and a flattening of the atomic radius, that accounts for the directionality of the interactions [55,56]. Thus, the halogen has an amphoteric character with an electropositive halogen bond ability along the σ -hole (C–X...O/N/S, $a \approx 180^\circ$) and an electronegative hydrogen bond acceptor perpendicular to the C–X bond (C–X...H, $a \approx 90^\circ$) [57–60]. Such halogen bonds have been detrimental in the understanding interactions of organic halogens in biological systems [61–65]. In our case, for compound **15**, I4 interacts with H4 ($d = 3.521 \text{ \AA}$, $a = 93.77^\circ$) and I4 also interact with the oxygen of the carbonyl of the acetate at C6 (I4...O, $d = 3.147 \text{ \AA}$; $a = 179.70^\circ$) (Figure 4b and Table 7). This result is in line with hydrogen (C–I...H) and halogen (C–I...O) interactions that show remarkable differences in term of geometrical features [66]. It is important to point out that similar interactions are also present in the packing of compound **13** (Cl4...H4 ($d = 3.625 \text{ \AA}$, $a = 96.94^\circ$) and Cl4...O ($d = 3.203 \text{ \AA}$, $a = 174.03^\circ$)) and compound **14** (Br4...H4 ($d = 3.526 \text{ \AA}$, $a = 95.15^\circ$) and Br4...O ($d = 3.143 \text{ \AA}$, $a = 177.49^\circ$)) (Table 7) (and see Supporting Information File 1). To the best of our knowledge, this is the first application of halogen bonding in the context of solid-state ordering of pyran inter-halides.




Table 7: Intermolecular X...H bond distances and angles for compound **13–15**.


Entry	Compound	D–H...A	<i>d</i> (D–H) (Å)	<i>d</i> (H...A) (Å)	<i>d</i> (D...A) (Å)	<i>a</i> (D–H–A) (°)	<i>a</i> (C–X–H) (°)
1	13	F2...H3	0.980	3.704	3.826	89.69	na
2		F2...H4	0.980	2.271	3.222	163.14	na
3		F3...H3	0.980	4.239	4.561	103.01	na
4		F3...H4	0.980	2.867	3.614	133.69	na
5		F3...H5	0.980	3.074	3.490	107.03	na
6		Cl4...H3	0.980	3.437	4.232	139.73	133.71
7		Cl4...H4	0.980	3.625	4.298	128.02	96.94
8		Cl4...H5	0.980	3.953	4.624	128.19	103.45
9	14	F2...H3	1.000	3.771	3.869	88.12	na
10		F2...H4	1.000	2.270	3.233	161.19	na
11		F3...H3	1.000	4.290	4.599	101.79	na
12		F3...H4	1.000	2.849	3.626	135.07	na
13		F3...H5	1.000	3.130	3.533	105.61	na
14		Br4...H3	1.000	3.316	4.132	139.90	133.33
15		Br4...H4	1.000	3.526	4.213	127.67	95.15
16		Br4...H5	1.000	3.887	4.564	127.36	102.99
17	15	F2...H3	0.980	3.867	3.950	87.64	na
18		F2...H4	0.981	2.356	3.293	159.70	na
19		F3...H3	0.980	4.382	4.673	101.24	na
20		F3...H4	0.981	2.886	3.668	137.42	na
21		F3...H5	0.980	3.248	3.612	103.92	na
22		I4...H3	0.980	3.285	4.086	140.25	132.28
23		I4...H4	0.981	3.521	4.192	127.59	93.77
24		I4...H5	0.980	3.905	4.559	126.76	102.70

Our interest in the synthesis and conformation of multivincinal inter-halides motivated us to use density functional theory (DFT) calculations to understand the preference of talose analogues to adopt 4C_1 -like conformations. DFT computations were performed using Gaussian 16 revision B.01 [67] with the CAM-B3LYP [68–70] functional and the Def2TZVP basis set [71], which includes effective core potentials for iodine. Empirical dispersion was accounted with Grimme's D3 [72,73] correction including Becke–Johnson damping [74]. Computations were performed both in the gas phase (i.e., individual molecules with thermal corrections at 298.15 K based on ideal gas assumptions) and in a chloroform solution, using the polarizable continuum model (PCM) [75]. A natural bonding orbital (NBO) analysis was performed to study the effects of hyperconjugation from C–F antibonding orbitals [76].

First, dipole moments, enthalpy and Gibbs free energy differences between 1C_4 and 4C_1 chair structures are shown in Table 8. In the gas phase, there is little difference in enthalpy

between the two structures. For Cl, Br, and I, the 1C_4 structures are predicted to have slightly lower enthalpies. However, as these differences are much smaller than 1 kcal/mol, one should conclude that the structures are nearly degenerate. In solution, the picture is much clearer: the 4C_1 structure is always lower in enthalpy and Gibbs free energy, which corresponds with the experimental measurements. One can see that the gap between the two structures tends to decrease as the halogen becomes larger (the minor exception being bromine). Dipole moments for the 4C_1 structures are 2 Debye larger than for 1C_4 structures, making them substantially more stable in chloroform.

Both chair structures have multiple C–F bonds in gauche arrangements which are stable due to hyperconjugation: there is donation from C–H bonding orbitals to C–F (or C–X) antibonding orbitals that are aligned with one another. The principal difference between the two structures is that in the 1C_4 structure there are two such interactions whereas in the 4C_1

Table 8: Dipole moments (Debye), enthalpy and Gibbs free energy differences (kcal/mol) computed (CAM-B3LYP-D3BJ/Def2TZVP) for 1C_4 and 4C_1 chair structures in the gas phase and in chloroform (PCM). Thermal corrections are reported at 298.15 K.


Entry	X	Gas phase				Chloroform			
		μ 1C_4	μ 4C_1	ΔH	ΔG	μ 1C_4	μ 4C_1	ΔH	ΔG
1	F	4.55	6.48	0.98	2.51	5.95	8.09	2.47	3.90
2	Cl	4.45	6.34	−0.03	1.31	5.93	7.93	1.29	2.21
3	Br	4.40	6.30	−0.13	1.18	5.89	7.94	1.19	2.55
4	I	4.32	6.07	−0.03	1.17	5.78	7.64	1.04	1.86

structure there are three. One can see that in the 4C_1 structure the C5–H5 bond will donate to the antibonding orbital of the C4–X4 bond, and the C3–H3 bond will donate to the antibonding orbitals of the C4–X4 and the C2–F2 bonds. In the 1C_4 structure, the C4–H4 and C2–H2 bonds will both donate to the antibonding orbital of the C3–F3 bond. These effects can be shown with an NBO analysis: the Kohn–Sham orbitals produced from DFT are localized to describe the system as one dominant resonance structure. As the Fock matrix is not diagonal in terms of the NBOs, coupling between orbitals can be quantified with second order perturbation theory. These

couplings represent donation from an occupied NBO to an unoccupied NBO that would stabilize the system. The results are presented in Table 9. In both cases, there is also donation from the halogen lone pairs to C–H and C–C antibonding orbitals, but as these effects were near equivalent in both chair structures, they are omitted.

Conclusion

We described the synthesis and conformational analysis of halogenated pyran analogues of D-talose. All analogues adopt standard 4C_1 -like conformations both in solution and in the solid-

Table 9: Second order perturbation theory energies of the Fock matrix in the NBO basis for 4C_1 and 1C_4 structures, from CAM-B3LYP-D3BJ/Def2TZVP results. Only results from CHCl₃ solvation (PCM) are reported as gas phase results showed no qualitative difference. Entries marked – are below the 0.50 kcal/mol threshold.


Entry	Donor σ	Acceptor σ^*	4C_1				1C_4			
			X = F	X = Cl	X = Br	X = I	X = F	X = Cl	X = Br	X = I
1	C5–H5	C4–X4	5.07	6.58	7.46	7.83	–	–	–	–
2	C4–H4	C4–X4	1.05	0.88	0.97	0.75	1.10	0.80	0.81	0.56
3	C4–H4	C3–F3	–	–	–	–	5.18	5.53	5.67	5.88
4	C3–H3	C4–X4	5.36	6.59	7.46	7.71	0.61	–	–	–
5	C3–H3	C3–F3	1.05	1.03	1.02	1.02	1.00	1.02	1.02	1.05
6	C3–H3	C2–F2	5.00	5.18	5.20	5.22	0.58	0.59	0.59	0.61
7	C2–H2	C3–F3	0.50	0.50	0.50	0.52	4.63	4.61	4.58	4.58
8	C2–H2	C2–F2	1.01	0.99	0.99	0.99	1.01	1.00	1.00	1.01
9	C1–H1	C2–F2	1.10	1.21	1.22	1.26	0.53	0.56	0.57	0.56

state. The conformations were corroborated using DFT calculations by looking at the energy, enthalpy and Gibbs' free energy differences between 1C_4 and 4C_1 chair structures. Crystallographic data of halogenated analogues shows intra-annular torsion angles demonstrated with the increasing distance between F2...X4 in relation with the nature of the halogen at C4: F ($d = 2.82 \text{ \AA}$) < Cl ($d = 3.06 \text{ \AA}$) < Br ($d = 3.14 \text{ \AA}$) < I ($d = 3.23 \text{ \AA}$). Moreover, the Cremer–Pople ring puckering parameters show suitable differences in the distortion of the chair conformations. Crystal packing arrangements showed that the halogen at C4 contributed in the nonbonding (along the σ -hole) and hydrogen bond (perpendicular to the C–X bond) interactions. Finally, this study should be of general interest in the understanding of weak interactions that are now important to so many areas of chemistry, such as crystal engineering and supramolecular chemistry.

Supporting Information

Supporting Information File 1

Experimental and analytical data, crystal structure determination and NMR spectra.

[<https://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-20-208-S1.pdf>]

Funding

This work was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) and Université Laval. O. L. thanks NSERC for a postgraduate fellowship. P. A. J. acknowledges funding from NSERC. This research was enabled by the Digital Research Alliance of Canada.

Author Contributions

Olivier Lessard: conceptualization; data curation; formal analysis; investigation; visualization; writing – original draft; writing – review & editing. Mathilde Grosset-Magagne: data curation; investigation; validation. Paul A. Johnson: resources; software; writing – original draft; writing – review & editing. Denis Giguère: conceptualization; funding acquisition; project administration; resources; supervision; visualization; writing – original draft; writing – review & editing.

ORCID® iDs

Paul A. Johnson - <https://orcid.org/0000-0003-1133-6880>

Denis Giguère - <https://orcid.org/0000-0003-2209-1428>

Data Availability Statement

All data that supports the findings of this study is available in the published article and/or the supporting information to this article.

Preprint

A non-peer-reviewed version of this article has been previously published as a preprint: <https://doi.org/10.3762/bxiv.2024.37.v1>

References

- Zhou, Y.; Wang, J.; Gu, Z.; Wang, S.; Zhu, W.; Aceña, J. L.; Soloshonok, V. A.; Izawa, K.; Liu, H. *Chem. Rev.* **2016**, *116*, 422–518. doi:10.1021/acs.chemrev.5b00392
- Wang, J.; Sánchez-Roselló, M.; Aceña, J. L.; del Pozo, C.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. *Chem. Rev.* **2014**, *114*, 2432–2506. doi:10.1021/cr4002879
- O'Hagan, D. *J. Fluorine Chem.* **2010**, *131*, 1071–1081. doi:10.1016/j.jfluchem.2010.03.003
- Gillis, E. P.; Eastman, K. J.; Hill, M. D.; Donnelly, D. J.; Meanwell, N. A. *J. Med. Chem.* **2015**, *58*, 8315–8359. doi:10.1021/acs.jmedchem.5b00258
- Meanwell, N. A. *J. Med. Chem.* **2018**, *61*, 5822–5880. doi:10.1021/acs.jmedchem.7b01788
- Linclau, B.; Ardá, A.; Reichardt, N.-C.; Sollogoub, M.; Unione, L.; Vincent, S. P.; Jiménez-Barbero, J. *Chem. Soc. Rev.* **2020**, *49*, 3863–3888. doi:10.1039/c9cs00099b
- Huonnic, K.; Linclau, B. *Chem. Rev.* **2022**, *122*, 15503–15602. doi:10.1021/acs.chemrev.2c00086
- Lessard, O.; Lainé, D.; Giguère, D. *Eur. J. Org. Chem.* **2024**, *27*, e202400120. doi:10.1002/ejoc.202400120
- Govindaraj, V.; Ungati, H.; Jakka, S. R.; Bose, S.; Mughesh, G. *Chem. – Eur. J.* **2019**, *25*, 11180–11192. doi:10.1002/chem.201902243
- Ungati, H.; Govindaraj, V.; Nair, C. R.; Mughesh, G. *Chem. – Eur. J.* **2019**, *25*, 3391–3399. doi:10.1002/chem.201806122
- Gerebtzoff, G.; Li-Blatter, X.; Fischer, H.; Frentzel, A.; Seelig, A. *ChemBioChem* **2004**, *5*, 676–684. doi:10.1002/cbic.200400017
- Nunes, R. S.; Vila-Vigosa, D.; Costa, P. J. *J. Am. Chem. Soc.* **2021**, *143*, 4253–4267. doi:10.1021/jacs.0c12470
- Heidrich, J.; Sperl, L. E.; Boeckler, F. M. *Front. Chem. (Lausanne, Switz.)* **2019**, *7*, 9. doi:10.3389/fchem.2019.00009
- Xu, Z.; Yang, Z.; Liu, Y.; Lu, Y.; Chen, K.; Zhu, W. *J. Chem. Inf. Model.* **2014**, *54*, 69–78. doi:10.1021/ci400539q
- Mendez, L.; Henriquez, G.; Sirimulla, S.; Narayan, M. *Molecules* **2017**, *22*, 1397. doi:10.3390/molecules22091397
- Hernandes, M. Z.; Cavalcanti, S. M. T.; Moreira, D. R. M.; de Azevedo Junior, W. F.; Leite, A. C. L. *Curr. Drug Targets* **2010**, *11*, 303–314. doi:10.2174/138945010790711996
- Wilcken, R.; Zimmermann, M. O.; Lange, A.; Joerger, A. C.; Boeckler, F. M. *J. Med. Chem.* **2013**, *56*, 1363–1388. doi:10.1021/jm3012068
- Chiodi, D.; Ishihara, Y. *J. Med. Chem.* **2023**, *66*, 5305–5331. doi:10.1021/acs.jmedchem.2c02015
- Tan, Y.; Luo, S.; Li, D.; Zhang, N.; Jia, S.; Liu, Y.; Qin, W.; Song, C. E.; Yan, H. *J. Am. Chem. Soc.* **2017**, *139*, 6431–6436. doi:10.1021/jacs.7b02076
- Bucher, C.; Deans, R. M.; Burns, N. Z. *J. Am. Chem. Soc.* **2015**, *137*, 12784–12787. doi:10.1021/jacs.5b08398
- Gribble, G. W. *Mar. Drugs* **2015**, *13*, 4044–4136. doi:10.3390/md13074044
- Gál, B.; Bucher, C.; Burns, N. Z. *Mar. Drugs* **2016**, *14*, 206. doi:10.3390/md14110206
- Lessard, O.; Lainé, D.; Fecteau, C.-É.; Johnson, P. A.; Giguère, D. *Org. Chem. Front.* **2022**, *9*, 6566–6572. doi:10.1039/d2qo01433e

24. Denavit, V.; Lainé, D.; St-Gelais, J.; Johnson, P. A.; Giguère, D. *Nat. Commun.* **2018**, *9*, 4721. doi:10.1038/s41467-018-06901-y
25. Denavit, V.; St-Gelais, J.; Tremblay, T.; Giguère, D. *Chem. – Eur. J.* **2019**, *25*, 9272–9279. doi:10.1002/chem.201901346
26. Denavit, V.; Lainé, D.; Bouzriba, C.; Shanina, E.; Gillon, É.; Fortin, S.; Rademacher, C.; Imbert, A.; Giguère, D. *Chem. – Eur. J.* **2019**, *25*, 4478–4490. doi:10.1002/chem.201806197
27. St-Gelais, J.; Bouchard, M.; Denavit, V.; Giguère, D. *J. Org. Chem.* **2019**, *84*, 8509–8522. doi:10.1021/acs.joc.9b00795
28. Lainé, D.; Denavit, V.; Lessard, O.; Carrier, L.; Fecteau, C.-É.; Johnson, P. A.; Giguère, D. *Beilstein J. Org. Chem.* **2020**, *16*, 2880–2887. doi:10.3762/bjoc.16.237
29. St-Gelais, J.; Côté, É.; Lainé, D.; Johnson, P. A.; Giguère, D. *Chem. – Eur. J.* **2020**, *26*, 13499–13506. doi:10.1002/chem.202002825
30. Leclerc, C.; St-Gelais, J.; Cecioni, S.; Giguère, D. *J. Fluorine Chem.* **2024**, *273*, 110232. doi:10.1016/j.jfluchem.2023.110232
31. Lainé, D.; Lessard, O.; St-Gelais, J.; Giguère, D. *Chem. – Eur. J.* **2021**, *27*, 3799–3805. doi:10.1002/chem.202004646
32. Passaglia, L.; Zanardi, M. M.; Sarotti, A. M. *Org. Biomol. Chem.* **2024**, *22*, 2435–2442. doi:10.1039/d3ob02077k
33. Deposition number 2344127 (for **13**), 2344124 (for **14**), and 2344126 (for **15**) contains the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service.
34. Deposition number 1824902 (for **17**) contains the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service.
35. Hansen, L. K.; Hordvik, A. *Acta Chem. Scand., Ser. A* **1977**, *31*, 187–191. doi:10.3891/acta.chem.scand.31a-0187
36. Hunter, L.; Slawin, A. M. Z.; Kirsch, P.; O'Hagan, D. *Angew. Chem., Int. Ed.* **2007**, *46*, 7887–7890. doi:10.1002/anie.200701988
37. Hunter, L.; O'Hagan, D.; Slawin, A. M. Z. *J. Am. Chem. Soc.* **2006**, *128*, 16422–16423. doi:10.1021/ja066188p
38. Bentler, P.; Erdeljac, N.; Bussmann, K.; Ahlqvist, M.; Knerr, L.; Bergander, K.; Daniluc, C. G.; Gilmour, R. *Org. Lett.* **2019**, *21*, 7741–7745. doi:10.1021/acs.orglett.9b02662
39. O'Hagan, D. *Chem. – Eur. J.* **2020**, *26*, 7981–7997. doi:10.1002/chem.202000178
40. Marchand, A. P.; Sorokin, V. D.; Rajagopal, D.; Bott, S. G. *Tetrahedron* **1994**, *50*, 9933–9942. doi:10.1016/s0040-4020(01)89608-3
41. Rodríguez-Vázquez, N.; Salzinger, S.; Silva, L. F.; Amorín, M.; Granja, J. R. *Eur. J. Org. Chem.* **2013**, 3477–3493. doi:10.1002/ejoc.201201565
42. Jeffrey, G. A. *Acta Crystallogr., Sect. B: Struct. Sci.* **1990**, *46*, 89–103. doi:10.1107/s0108768189012449
43. Bock, K.; Duus, J. Ø. *J. Carbohydr. Chem.* **1994**, *13*, 513–543. doi:10.1080/07328309408011662
44. Linclau, B.; Golten, S.; Light, M.; Sebban, M.; Oulyadi, H. *Carbohydr. Res.* **2011**, *346*, 1129–1139. doi:10.1016/j.carres.2011.04.007
45. Kim, H. W.; Rossi, P.; Shoemaker, R. K.; DiMaggio, S. G. *J. Am. Chem. Soc.* **1998**, *120*, 9082–9083. doi:10.1021/ja9803714
46. Allen, F. H.; Kennard, O.; Watson, D. G.; Brammer, L.; Orpen, A. G.; Taylor, R. J. *Chem. Soc., Perkin Trans. 2* **1987**, S1–S19. doi:10.1039/p2987000000s1
47. Jeffrey, G. A.; Pople, J. A.; Radom, L. *Carbohydr. Res.* **1972**, *25*, 117–131. doi:10.1016/s0008-6215(00)82752-4
48. O'Hagan, D. *Chem. Soc. Rev.* **2008**, *37*, 308–319. doi:10.1039/b711844a
49. Bondi, A. *J. Phys. Chem.* **1964**, *68*, 441–451. doi:10.1021/j100785a001
50. Cremer, D.; Pople, J. A. *J. Am. Chem. Soc.* **1975**, *97*, 1354–1358. doi:10.1021/ja00839a011
51. Jeffrey, G. A.; Yates, J. H. *Carbohydr. Res.* **1979**, *74*, 319–322. doi:10.1016/s0008-6215(00)84786-2
52. Steiner, T. *Angew. Chem., Int. Ed.* **2002**, *41*, 48–76. doi:10.1002/1521-3773(20020104)41:1<48::aid-anie48>3.0.co;2-u
53. Bresciani, S.; Lebl, T.; Slawin, A. M. Z.; O'Hagan, D. *Chem. Commun.* **2010**, *46*, 5434–5436. doi:10.1039/c0cc01128b
54. O'Hagan, D. *Chem. Rec.* **2023**, *23*, e202300027. doi:10.1002/tcr.202300027
55. Clark, T.; Hennemann, M.; Murray, J. S.; Politzer, P. J. *Mol. Model.* **2007**, *13*, 291–296. doi:10.1007/s00894-006-0130-2
56. Politzer, P.; Murray, J. S.; Clark, T. *Phys. Chem. Chem. Phys.* **2013**, *15*, 11178–11189. doi:10.1039/c3cp00054k
57. Scholfield, M. R.; Zanden, C. M. V.; Carter, M.; Ho, P. S. *Protein Sci.* **2013**, *22*, 139–152. doi:10.1002/pro.2201
58. Nelyubina, Y. V.; Antipin, M. Y.; Dunin, D. S.; Kotov, V. Y.; Lyssenko, K. A. *Chem. Commun.* **2010**, *46*, 5325–5327. doi:10.1039/c0cc01094d
59. Brammer, L.; Bruton, E. A.; Sherwood, P. *Cryst. Growth Des.* **2001**, *1*, 277–290. doi:10.1021/cg015522k
60. Aakeröy, C. B.; Desper, J.; Helfrich, B. A.; Metrangola, P.; Pilati, T.; Resnati, G.; Stevenazzi, A. *Chem. Commun.* **2007**, *43*, 4236–4238. doi:10.1039/b707458a
61. Scholfield, M. R.; Ford, M. C.; Carlsson, A.-C. C.; Butta, H.; Mehl, R. A.; Ho, P. S. *Biochemistry* **2017**, *56*, 2794–2802. doi:10.1021/acs.biochem.7b00022
62. Lu, Y.; Wang, Y.; Zhu, W. *Phys. Chem. Chem. Phys.* **2010**, *12*, 4543–4551. doi:10.1039/b926326h
63. Auffinger, P.; Hays, F. A.; Westhof, E.; Ho, P. S. *Proc. Natl. Acad. Sci. U. S. A.* **2004**, *101*, 16789–16794. doi:10.1073/pnas.0407607101
64. Lin, F.-Y.; MacKerell, A. D., Jr. *J. Phys. Chem. B* **2017**, *121*, 6813–6821. doi:10.1021/acs.jpcc.7b04198
65. Lu, Y.; Wang, Y.; Xu, Z.; Yan, X.; Luo, X.; Jiang, H.; Zhu, W. *J. Phys. Chem. B* **2009**, *113*, 12615–12621. doi:10.1021/jp906352e
66. Decato, D. A.; Sun, J.; Boller, M. R.; Berryman, O. B. *Chem. Sci.* **2022**, *13*, 11156–11162. doi:10.1039/d2sc03792k
67. *Gaussian 16*, Revision B.01; Gaussian, Inc.: Wallingford CT, 2016.
68. Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648–5652. doi:10.1063/1.464913
69. Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785–789. doi:10.1103/physrevb.37.785
70. Yanai, T.; Tew, D. P.; Handy, N. C. *Chem. Phys. Lett.* **2004**, *393*, 51–57. doi:10.1016/j.cplett.2004.06.011
71. Weigend, F.; Ahlrichs, R. *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297–3305. doi:10.1039/b508541a
72. Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. *J. Chem. Phys.* **2010**, *132*, 154104. doi:10.1063/1.3382344
73. Grimme, S.; Ehrlich, S.; Goerigk, L. *J. Comput. Chem.* **2011**, *32*, 1456–1465. doi:10.1002/jcc.21759
74. Becke, A. D.; Johnson, E. R. *J. Chem. Phys.* **2006**, *124*, 014104. doi:10.1063/1.2139668
75. Tomasi, J.; Mennucci, B.; Cammi, R. *Chem. Rev.* **2005**, *105*, 2999–3094. doi:10.1021/cr9904009
76. Foster, J. P.; Weinhold, F. *J. Am. Chem. Soc.* **1980**, *102*, 7211–7218. doi:10.1021/ja00544a007

License and Terms

This is an open access article licensed under the terms of the Beilstein-Institut Open Access License Agreement (<https://www.beilstein-journals.org/bjoc/terms>), which is identical to the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0>). The reuse of material under this license requires that the author(s), source and license are credited. Third-party material in this article could be subject to other licenses (typically indicated in the credit line), and in this case, users are required to obtain permission from the license holder to reuse the material.

The definitive version of this article is the electronic one which can be found at:
<https://doi.org/10.3762/bjoc.20.208>