Supporting Information

for

Bis(β -lactosyl)-[60]fullerene as novel class of glycolipids useful for detection and decontamination of biological toxins of the *Ricinus communis* family

Hirofumi Dohi^{1*}, Takeru Kanazawa¹, Akihiro Saito², Keita Sato³, Hirotaka Uzawa⁴, Yasuo Seto³, Yoshihiro Nishida^{1*}

Address: ¹Department of Nanobiology, Graduate School of Advanced Integration Science, Chiba University, 1-33 Yayoi-cho, Inage-ku, Chiba 263–8522, Japan, ²Department of Materials and Life Science, Shizuoka Institute of Science and Technology, 2200-2 Toyosawa, Fukuroi, Shizuoka 437-8555, Japan, ³National Research Institute of Police Science, 6-3-1 Kashiwanoha, Kashiwa, Chiba 277-0882, Japan and ⁴Nanosystem Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), 1-1-1 Higashi, Tsukuba, 305-8565, Japan

Email: Hirofumi Dohi - <u>hdohi@faculty.chiba-u.jp</u>; Yoshihiro Nishida -YNishida@faculty.chiba-u.jp

*Corresponding author

Copies of ¹H and ¹³C NMR spectra for compounds 2, 3 and 4.



Figure S1. ¹H and ¹³C NMR spectra of compound 2.



Figure S2. ¹H and ¹³C NMR spectra of compound 3.



Figure S3. ¹H and ¹³C NMR spectra of compound 4.