

Supporting Information File 2

for

Impact of multivalent charge presentation on peptide– nanoparticle aggregation

Daniel Schöne¹, Boris Schade², Christoph Böttcher² and Beate Kokschr*¹

Address: ¹Institute of Chemistry and Biochemistry - Organic Chemistry, Freie Universität Berlin,
Takustr. 3, 14195 Berlin, Germany and ²Electron Microscopy, Freie Universität Berlin,
Fabeckstr. 36a, 14195 Berlin, Germany

Email: Beate Kokschr* - beate.kokschr@fu-berlin.de

*Corresponding author

TEM images




Figure S5: TEM of 100 μ M R1A3 in 10 mM Tris/HCl buffer pH 9. Sample was negative stained with 2 % PTA, defocus - 0.5 μ m.




Figure S6: TEM of 100 μ M R2A3 in 10 mM Tris/HCl buffer pH 9. Sample was negative stained with 2 % PTA, defocus - 0.5 μ m.




Figure S7: TEM of 100 μ M R2A4 in 10 mM Tris/HCl buffer pH 9. Sample was negative stained with 2 % PTA, defocus - 0.5 μ m.




Figure S8: TEM of 100 μ M R2A5 in 10 mM Tris/HCl buffer pH 9. Sample was negative stained with 2 % PTA, defocus - 0.5 μ m.




Figure S9: Stereo Cryo TEM of 100 μM R2A2 and 0.4 μM Au/MUA nanoparticles in 10 mM Tris/HCl buffer pH 9.