

Supporting Information

for

Ni nanoparticles on RGO as reusable heterogeneous catalyst: effect of Ni particle size and intermediate composite structures in C–S cross-coupling reaction

Debasish Sengupta^{†1}, Koushik Bhowmik^{‡2}, Goutam De^{*2} and Basudeb Basu^{*1}

Address: ¹Department of Chemistry, University of North Bengal, Darjeeling 734013, India. Fax: +91 353 2699001; Tel: +91 353 2776381 and ²Nano-Structured Materials Division, CSIR–Central Glass & Ceramic Research Institute, 196, Raja S. C. Mullick Road, Jadavpur, Kolkata 700032, India. Fax: +91 33 24730957; Tel: +91 33 23223403.

Email: Goutam De^{*} - gde@cgcri.res.in; Basudeb Basu^{*} - basu_nbu@hotmail.com,

^{*}Corresponding author

[‡]These authors have contributed equally

¹H and ¹³C NMR spectra (scanned) for compounds 3a–p

Table 2; entry 1





Table 2; entry 2





Table 2; entry 3




Table 2; entry 4





Table 2; entry 5





Table 2; entry 6





Table 2; entry 7





Table 2; entry 8





Table 2; entry 9





Table 2; entry 10





Table 2; entry 11





Table 2; entry 12





Table 2; entry 14





Table 2; entry 15





Table 2; entry 16




Table 2; entry 17

