

Supporting Information

for

Fast and efficient synthesis of microporous polymer nanomembranes via light-induced click reaction

Qi An¹, Youssef Hassan^{1,2}, Xiaotong Yan¹, Peter Krolla-Sidenstein¹, Tawheed Mohammed^{1,3}, Mathias Lang⁴, Stefan Bräse^{4,5} and Manuel Tsotsalas^{*1,4}

Address: ¹Institute of Functional Interfaces (IFG), Karlsruhe Institute of Technology (KIT), Hermann-von-Helmholtz-Platz 1, D-76344 Eggenstein-Leopoldshafen, Germany, ²Zewail City of Science and Technology, Center for Materials Science, Sheikh Zayed District, 6th of October City, 12588, Giza, Egypt, ³Institute of Physics and Technology, International X-ray Optics Lab, National Research Tomsk Polytechnic University (TPU), 30 Lenin ave., Tomsk 634050, Russia, ⁴Institute for Organic Chemistry (IOC), Karlsruhe Institute of Technology (KIT), Fritz-Haber-Weg 6, D-76131 Karlsruhe, Germany, and ⁵Institute of Toxicology and Genetics (ITG), Karlsruhe Institute of Technology (KIT), Hermann-von-Helmholtz-Platz 1, D-76344 Eggenstein-Leopoldshafen, Germany

Email: Manuel Tsotsalas - manuel.tsotsalas@kit.edu

* Corresponding author

Additional IRRA spectra




Figure S1: IRRA spectrum of Tetrakis(4-ethynyl-phenyl)methane (TPM-alkyne).




Figure S2: IRRA spectrum of Tetrakis(4-thiolphenyl)-methane (TPM-SH).




Figure S3: IRRA spectrum of 2-hydroxy-4'-(2-hydroxyethoxy)-2-methylpropiophenone.