



## Supporting Information


for



### **A chemoenzymatic synthesis of ceramide trafficking inhibitor HPA-12**


Seema V. Kanojia, Sucheta Chatterjee, Subrata Chattopadhyay and Dibakar Goswami



*Beilstein J. Org. Chem.* **2019**, *15*, 490–496. [doi:10.3762/bjoc.15.42](https://doi.org/10.3762/bjoc.15.42)


## NMR spectra



























8.298  
8.280  
8.243  
8.226  
7.440  
7.426  
7.398  
7.384  
7.369  
7.340  
7.326  
7.312  
7.264  
6.090  
6.076  
6.063  
5.818  
5.804  
5.798  
5.784  
5.770  
5.756  
5.750  
5.736  
5.165  
5.131  
5.103  
5.083


2.872  
2.857  
2.842  
2.828  
2.813  
2.759  
2.747  
2.733  
2.718  
2.706





(S)-4a






























































— 166.178  
 { 143.272  
 { 133.321  
 { 132.126  
 { 129.780  
 { 129.444  
 { 128.825  
 { 128.490  
 { 128.189  
 { 125.895  
 { 77.252  
 { 76.998  
 { 76.744  
 { 71.903  
 { 67.002  
 — 58.714  
 — 39.631

