



Supporting Information

for

A cyclopeptide and three oligomycin-class polyketides produced by an underexplored actinomycete of the genus *Pseudosporangium*

Shun Saito, Kota Atsumi, Tao Zhou, Keisuke Fukaya, Daisuke Urabe, Naoya Oku, Md. Rokon Ul Karim, Hisayuki Komaki and Yasuhiro Igarashi

Beilstein J. Org. Chem. **2020**, *16*, 1100–1110. doi:10.3762/bjoc.16.97

Copies of NMR spectra for compounds 1–4

Table of contents

Figure S1. ^1H NMR spectrum of pseudosporamide (1) (500 MHz, $\text{DMSO-}d_6$)	S3
Figure S2. ^{13}C NMR spectrum of 1 (125 MHz, $\text{DMSO-}d_6$)	S4
Figure S3. DEPT135 spectrum of 1 (125 MHz, $\text{DMSO-}d_6$)	S5
Figure S4. $^1\text{H-}^1\text{H}$ COSY spectrum of 1 (500 MHz, $\text{DMSO-}d_6$)	S6
Figure S5. HSQC spectrum of 1 (500 MHz, $\text{DMSO-}d_6$)	S7
Figure S6. HMBC spectrum of 1 (500 MHz, $\text{DMSO-}d_6$)	S8
Figure S7. ROESY spectrum of 1 (500 MHz, $\text{DMSO-}d_6$)	S9
Figure S8. HPLC chromatogram of hydrolysate from 1 and comparison to standard amino acids, derivatized with Marfey's reagent (FDLA)	S10
Figure S9. ^1H NMR spectrum of 5a (500 MHz, $\text{DMSO-}d_6$)	S11
Figure S10. $^1\text{H-}^1\text{H}$ COSY spectrum of 5a (500 MHz, $\text{DMSO-}d_6$)	S12
Figure S11. ^1H NMR spectrum of 5b (500 MHz, $\text{DMSO-}d_6$)	S13
Figure S12. $^1\text{H-}^1\text{H}$ COSY spectrum of 5b (500 MHz, $\text{DMSO-}d_6$)	S14
Figure S13. ^1H NMR spectrum of pseudosporamicin A (2) (500 MHz, CDCl_3)	S15
Figure S14. ^{13}C NMR spectrum of 2 (125 MHz, CDCl_3)	S16
Figure S15. $^1\text{H-}^1\text{H}$ COSY spectrum of 2 (500 MHz, CDCl_3)	S17
Figure S16. HSQC spectrum of 2 (500 MHz, CDCl_3)	S18
Figure S17. HMBC spectrum of 2 (500 MHz, CDCl_3)	S19
Figure S18. NOESY spectrum of 2 (500 MHz, CDCl_3)	S20
Figure S19. ^1H NMR spectrum of pseudosporamicin B (3) (500 MHz, CDCl_3)	S21
Figure S20. ^{13}C NMR spectrum of 3 (125 MHz, CDCl_3)	S22
Figure S21. $^1\text{H-}^1\text{H}$ COSY spectrum of 3 (500 MHz, CDCl_3)	S23
Figure S22. HSQC spectrum of 3 (500 MHz, CDCl_3)	S24
Figure S23. HMBC spectrum of 3 (500 MHz, CDCl_3)	S25
Figure S24. NOESY spectrum of 3 (500 MHz, CDCl_3)	S26
Figure S25. ^1H NMR spectrum of pseudosporamicin C (4) (500 MHz, CDCl_3)	S27
Figure S26. ^{13}C NMR spectrum of 4 (125 MHz, CDCl_3)	S28
Figure S27. $^1\text{H-}^1\text{H}$ COSY spectrum of 4 (500 MHz, CDCl_3)	S29
Figure S28. HSQC spectrum of 4 (500 MHz, CDCl_3)	S30
Figure S29. HMBC spectrum of 4 (500 MHz, CDCl_3)	S31
Figure S30. NOESY spectrum of 4 (500 MHz, CDCl_3)	S32
Figure S31. COSY, key HMBC and NOESY correlations of 3 and 4	S33

Figure S1. ^1H NMR spectrum of pseudosporamide (**1**) (500 MHz, $\text{DMSO}-d_6$).




Figure S2. ^{13}C NMR spectrum of **1** (125 MHz, $\text{DMSO-}d_6$).




Figure S3. DEPT135 spectrum of **1** (125 MHz, DMSO- d_6).




Figure S4. ^1H - ^1H COSY spectrum of **1** (500 MHz, $\text{DMSO-}d_6$).




Figure S5. HSQC spectrum of **1** (500 MHz, DMSO- d_6).




Figure S6. HMBC spectrum of **1** (500 MHz, DMSO- d_6).





Figure S7. ROESY spectrum of **1** (500 MHz, DMSO- d_6).






Figure S8. HPLC chromatogram of hydrolysate from **1** and comparison to standard amino acids, derivatized with Marfey's reagent (FDLA).

Column: Cosmosil AR-II (Nacalai Tesque, 4.6×250 mm)

Solvent: MeCN-0.1% HCO₂H

Gradient MeCN (%): 0-30 min: 15-80%, 30-35 min: 80%

Flow rate: 1.0 mL/min

Absorbance: 340 nm

Figure S9. ^1H NMR spectrum of **5a** (500 MHz, $\text{DMSO}-d_6$).




Figure S10. ^1H - ^1H COSY spectrum of **5a** (500 MHz, $\text{DMSO-}d_6$).




Figure S11. ^1H NMR spectrum of **5b** (500 MHz, $\text{DMSO}-d_6$).




Figure S12. ^1H - ^1H COSY spectrum of **5b** (500 MHz, $\text{DMSO-}d_6$).




Figure S13. ^1H NMR spectrum of pseudosporamicin A (**2**) (500 MHz, CDCl_3).




Figure S14. ^{13}C NMR spectrum of **2** (125 MHz, CDCl_3).




Figure S15. ^1H - ^1H COSY spectrum of **2** (500 MHz, CDCl_3).




Figure S16. HSQC spectrum of **2** (500 MHz, CDCl₃).




Figure S17. HMBC spectrum of **2** (500 MHz, CDCl_3).




Figure S18. NOESY spectrum of **2** (500 MHz, CDCl_3).




Figure S20. ^{13}C NMR spectrum of **3** (125 MHz, CDCl_3).




Figure S21. ^1H - ^1H COSY spectrum of **3** (500 MHz, CDCl_3).




Figure S22. HSQC spectrum of **3** (500 MHz, CDCl₃).




Figure S23. HMBC spectrum of **3** (500 MHz, CDCl₃).




Figure S24. NOESY spectrum of **3** (500 MHz, CDCl₃).




Figure S25. ^1H NMR spectrum of pseudosporamicin C (**4**) (500 MHz, CDCl_3).




Figure S26. ^{13}C NMR spectrum of **4** (125 MHz, CDCl_3).




Figure S27. ^1H - ^1H COSY spectrum of **4** (500 MHz, CDCl_3).




Figure S28. HSQC spectrum of **4** (500 MHz, CDCl₃).




Figure S29. HMBC spectrum of **4** (500 MHz, CDCl₃).





Figure S30. NOESY spectrum of **4** (500 MHz, CDCl₃).



Chemical structures of compounds **3** and **4** are shown, illustrating the assignment of carbon and proton signals using COSY, HMBC, and NOESY correlations.

Legend:

- COSY
- HMBC
- NOESY