



Supporting Information

for

Synthetic approach to 2-alkyl-4-quinolones and 2-alkyl-4-quinolone-3-carboxamides based on common β -keto amide precursors


Yordanka Mollova-Sapundzhieva, Plamen Angelov, Danail Georgiev and Pavel Yanev

Beilstein J. Org. Chem. **2023**, *19*, 1804–1810. [doi:10.3762/bjoc.19.132](https://doi.org/10.3762/bjoc.19.132)

Processed NMR spectra



Table of contents	Page
Processed NMR spectra (¹ H/ ¹³ C)	
1-Hydroxy-2-alkyl-4-quinolone-3-carboxamides 4	S2
2-Alkyl-4-quinolone-3-carboxamides 5	S7
2-Alkyl-4-quinolones 8	S16

Primary NMR data available at: <https://doi.org/10.5281/zenodo.8337063>




4a



400 MHz, DMSO-*d*₆




4b





400 MHz, DMSO-*d*₆






4c



400 MHz, DMSO-*d*₆






4d



400 MHz, DMSO-*d*₆




4e





400 MHz, DMSO-*d*₆




5a





400 MHz, DMSO-*d*₆






5b



400 MHz, DMSO-*d*₆




5c





400 MHz, DMSO-d₆




5d





400 MHz, DMSO-*d*₆




5e





400 MHz, DMSO- d_6




5f





400 MHz, DMSO-*d*₆




5g





400 MHz, DMSO-*d*₆






5h



400 MHz, DMSO-*d*₆






5i



400 MHz, DMSO-*d*₆




8a





400 MHz, DMSO-d₆




8b





400 MHz, DMSO-*d*₆




8c



400 MHz, DMSO-*d*₆



8d



400 MHz, DMSO-*d*₆

