

Supporting Information

for

Synthesis, biological and electrochemical evaluation of glycidyl esters of phosphorus acids as potential anticancer drugs

Almaz A. Zagidullin, Emil R. Bulatov, Mikhail N. Khrizanforov, Damir R. Davletshin, Elvina M. Gilyazova, Ivan A. Strelkov and Vasily A. Miluykov

Beilstein J. Org. Chem. 2025, 21, 1909-1916. doi:10.3762/bjoc.21.148

¹H, ³¹P NMR and IR spectra of compounds 1–3

³¹P, ¹H NMR, IR spectra for diglycidyl methylphosphonate (1).




Figure S1. ³¹P{¹H} NMR spectrum of 1 in CDCl₃.




Figure S2. ¹H NMR spectrum of 1 in CDCl₃.




Figure S3. IR spectrum of 1 in CDCl₃.

³¹P, ¹H NMR, IR spectra for diglycidyl methylphosphate (2).




Figure S4. ³¹P{¹H} NMR spectrum of 2 in CDCl₃.




Figure S5. ¹H NMR spectrum of 2 in CDCl₃.




Figure S6. IR spectrum of 2.

³¹P, ¹H NMR, IR spectra for triglycidyl phosphate (3).




Figure S7. ³¹P{¹H} NMR spectrum of 3 in CDCl₃.




Figure S8. ¹H NMR spectrum of 3 in CDCl₃.




Figure S9. IR spectrum of 3.