



Supporting Information

for

A multicomponent reaction-initiated synthesis of imidazopyridine-fused isoquinolinones

Ashutosh Nath, John Mark Awad and Wei Zhang

Beilstein J. Org. Chem. **2025**, 21, 1161–1169. [doi:10.3762/bjoc.21.92](https://doi.org/10.3762/bjoc.21.92)

General reaction procedures, compound characterization data, and copies of NMR spectra

Table of Contents

General Information.....	S2
General procedure for the synthesis of GBB products 4	S2
General procedure of N-acylation for the synthesis of products 6	S2
General procedure for IMDA and dehydrative re-aromatization products 8	S2
X-ray crystallography report of 8a and 6t	S7
Analytical characterization data of products 6a and 8	S9
¹ H NMR, ¹³ C NMR spectra of products 6a and 8	S13

General Information.

All first reactions were conducted in a sealed Biotage microwave reaction vial containing 2–5 mL, unless stated otherwise. Analytical thin layer chromatography (TLC) was conducted on silica gel plates including an F-254 indicator, and chemicals were seen using UV light irradiation. Column chromatography was performed utilizing silica gel (200–300 mesh) under elevated pressure. The ^1H , and ^{13}C spectroscopic data were acquired using Bruker Mercury Plus 400 MHz or Bruker AVANCE NEO 500 MHz NMR spectrometers. Chemical shifts were expressed in parts per million (ppm) relative to internal TMS for ^1H NMR data and deuterated solvent for ^{13}C NMR data. ^1H NMR coupling constants were expressed in Hz, with multiplicity denoted as follows: s (singlet); d (doublet); t (triplet); q (quartet); m (multiplet); dd (doublet of doublets); and td (triplet of doublets). LC–MS were performed on an Agilent 2100 system with C_{18} column (5.0 μm , 6.0 $\times$ 50 mm). The mobile phases were ACN and H_2O both containing 0.05% trifluoroacetic acid. A linear gradient was used to increase from 25:75 (v/v) MeOH/ H_2O to 100% MeOH in 7.0 min at a flow rate of 0.7 mL/min. UV detections were conducted at 210 nm, 254 nm and 280 nm.

General procedure for the synthesis of GBB product 4

The initial GBB reactions for making imidazo[1,2-*a*] pyridines **4** were conducted using aminopyridines **1** (0.5 mmol), isocyanides **3** (0.6 mmol, 1.2 equiv), and furfuraldehyde **2** (0.6 mmol, 1.2 equiv) in 3:1 DCM/MeOH (4 mL) were conducted using $\text{Yb}(\text{OTf})_3$ (0.04 mmol, 0.08 equiv) as a Lewis acid catalyst under microwave irradiation at 100 °C for 1 h (Scheme 2, Table S1). Nineteen distinct adducts **4** were obtained in 89–98% yields.

General procedure of *N*-acylation for the synthesis of product 6

Reactions of **4** with acryloyl chloride **5** (1.5 equiv) in the presence of Et_3N (2 equiv) at room temperature in anhydrous CH_2Cl_2 for 6 h afforded 19 *N*-acylated compounds **6** in 80–90% yields (Table S2) [10]. Further purification was conducted by flash chromatography with 1:6 EtOAc/*n*-hexane. The adduct was confirmed by NMR.

General procedure for IMDA and dehydrative re-aromatization product 8

After step-2 isolated *N*-acylation product **6** and reflux in dichlorobenzene solvent 4 h at 180 °C with 0.08 equiv. Lewis's acid AlCl_3 (Table S3). During this reaction we checked crude LC–MS and observed DA-adduct **7** and ring open product **8**. After 4 h reaction isolated ring open product **8** and purified with Ethyl acetate/hexane (30:70). We confirmed product structure from ^1H , ^{13}C NMR and x-ray crystal structure analysis.

Table S1: Three-component GBB cycloaddition for the syntheses of **4**.





Table S2: *N*-Acylation of fused imidazo[1,2-*a*]pyridines.





Table S3: Substrate scope for the reaction of imidazopyridine-fused isoquinolinones **8**.



[a] Reactions of **6** were carried out using $AlCl_3$ (10 mol %) in 1,2-dichlorobenzene at 180 °C for 4 h.








Figure S1: LC-MS of overserving IMDA Adduct 7a.

X-ray crystallographic analysis of 8a (CCDC: 2429172)



Cell	a = 19.882(3) Å; b = 7.8944(9) Å; c = 20.396(3) Å	
	a = 90°; b = 90°; g = 90°.	
Temperature	300(2) K	
	Calculated	Reported
Volume	3201.3(7) Å ³	3201.3(7) Å ³
Space group	Pbca	Pbca
Moiety formula	C ₁₈ H ₁₆ BrN ₃ O	C ₁₈ H ₁₆ BrN ₃ O
Sum formula	C ₁₈ H ₁₆ BrN ₃ O	C ₁₈ H ₁₆ BrN ₃ O
Z	8	8
μ (mm ⁻¹)	2.576	2.576
F000	1504	1504
CCDC: 2429172		

X-ray crystallographic analysis of **6t** (CCDC: 2429579)




6t

Cell	a = 11.5629(12) Å; b = 11.0333(12) Å; c = 14.0321(16) Å	
Temperature	a = 90°; b = 106.650°; g = 90°. 300(2) K	
Volume	Calculated 1715.1(3) Å ³	Reported 1715.1(3) Å ³
Space group	P2 ₁ /n	P2 ₁ /n
Moiety formula	C ₁₈ H ₁₉ N ₃ OS	C ₁₈ H ₁₉ N ₃ OS
Sum formula	C ₁₈ H ₁₉ N ₃ OS	C ₁₈ H ₁₉ N ₃ OS
Z	4	4
μ (mm ⁻¹)	2.576	2.576
F000	1504	1504
CCDC: 2429579		


Analytical characterization data of products **6a** and **8**.

N-(2-(4-Bromofuran-2-yl)imidazo[1,2-*a*]pyridin-3-yl)-*N*-butylacrylamide (**6a**)




Light yellow solid (90% yield), Chemical Formula: C₁₈H₁₈BrN₃O₂. ¹H NMR (400 MHz, cdcl₃) δ 7.79 (d, *J* = 6.8 Hz, 1H), 7.63 (d, *J* = 9.1 Hz, 1H), 7.48 (s, 1H), 7.36 – 7.26 (m, 1H), 6.92 (t, *J* = 6.8 Hz, 1H), 6.81 (s, 1H), 6.43 (d, *J* = 16.7 Hz, 1H), 5.80 (dd, *J* = 16.7, 10.3 Hz, 1H), 5.50 (d, *J* = 10.4 Hz, 1H), 3.87 (ddd, *J* = 15.5, 10.1, 5.7 Hz, 1H), 3.67 (td, *J* = 13.3, 11.9, 5.9 Hz, 1H), 1.45 – 1.20 (m, 5H), 0.97 – 0.81 (m, 3H).

2-Bromo-6-butylpyrido[2',1':2,3]imidazo[4,5-*c*]isoquinolin-5(6*H*)-one (**8a**)




Yellow solid (85% yield), Chemical Formula: C₁₈H₁₆BrN₃O. ¹H NMR (400 MHz, cdcl₃) δ 8.57 (d, *J* = 1.9 Hz, 1H), 8.37 – 8.30 (m, 2H), 7.69 (d, *J* = 9.2 Hz, 1H), 7.63 (dd, *J* = 8.7, 1.9 Hz, 1H), 7.26 – 7.15 (m, 2H), 6.87 (t, *J* = 7.0 Hz, 1H), 4.63 (t, *J* = 7.9 Hz, 2H), 1.88 (q, *J* = 7.9 Hz, 2H), 1.59 – 1.50 (m, 2H), 1.02 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (100 MHz, cdcl₃) δ 158.5, 136.25, 134.2, 127.2, 124.1, 123.0, 122.6, 120.1, 118.1, 118.0, 116.2, 114.4, 112.0, 107.2, 41.2, 29.9, 17.9, 11.7. HRMS (ESI) calcd for C₁₈H₁₆BrN₃O *m/z*: 369.0477, found 369.0480.

6-Benzyl-2-bromopyrido[2',1':2,3]imidazo[4,5-*c*]isoquinolin-5(6*H*)-one (**8b**)




Light yellow solid (82% yield), C₂₁H₁₄BrN₃O. ¹H NMR (399 MHz,) δ 8.55 (d, *J* = 2.0 Hz, 1H), 8.40 – 8.31 (m, 5H), 7.70 – 7.61 (m, 4H), 7.17 (dd, *J* = 9.6, 1.8 Hz, 2H), 4.61 (t, *J* = 7.9 Hz, 2H). ¹³C NMR (100 MHz, cdcl₃) δ 150.1, 137.2, 130.3, 130.2, 128.9, 128.8, 126.3, 126.3, 125.1, 121.8, 120.8, 119.9, 105.1, 104.9, 46.6. HRMS (ESI) calcd for C₂₁H₁₄BrN₃O *m/z*: 403.0320, found 403.0341.

2-Bromo-6-isopropylpyrido[2',1':2,3]imidazo[4,5-*c*]isoquinolin-5(6*H*)-one (**8c**)




Off-white solid (77% yield), Chemical Formula: C₁₇H₁₄BrN₃O. ¹H NMR (399 MHz,) δ 8.57 (d, *J* = 1.9 Hz, 1H), 8.37 – 8.30 (m, 2H), 7.69 (d, *J* = 9.2 Hz, 1H), 7.63 (dd, *J* = 8.7, 1.9 Hz, 1H), 7.26 – 7.15 (m, 1H), 6.87 (t, *J* = 7.0 Hz, 1H), 5.09 – 4.75 (m, 1H), 1.89 – 1.83 (m, 6H). ¹³C NMR (100 MHz, cdcl₃) δ 168.3, 148.2, 131.9, 125.1, 124.1, 120.9, 120.1, 111.5, 110.2, 108.2, 106.3, 44.9, 20.1. HRMS (ESI) calcd for C₁₇H₁₄BrN₃O *m/z*: 355.0320, found 355.0624.

2-Bromo-6-cyclohexylpyrido[2',1':2,3]imidazo[4,5-*c*]isoquinolin-5(6*H*)-one (**8d**)




Off-white solid (70% yield), Chemical Formula: C₂₀H₁₈BrN₃O. ¹H NMR (399 MHz, cdcl₃) δ 8.38 – 8.31 (m, 1H), 8.26 (d, *J* = 8.5 Hz, 0H), 8.14 (d, *J* = 7.4 Hz, 1H), 7.69 (d, *J* = 9.3 Hz, 1H), 7.50 – 7.43 (m, 1H), 7.18 (s, 0H), 6.89 (t, *J* = 7.0 Hz, 1H), 4.39 (d, *J* = 12.1 Hz, 1H), 2.95 – 2.85 (m, 2H), 1.99 (dd, *J* = 26.2, 11.4 Hz, 5H), 0.85 (d, *J* = 17.3 Hz, 3H). ¹³C NMR (100 MHz, cdcl₃) δ 167.4, 143.1, 135.9, 135.2, 130.6, 128.0, 126.6, 126.3, 125.9, 124.9, 121.6, 117.9, 113.4, 48.2, 30.6, 20.3, 13.7.

2-Bromo-6-(2-morpholinoethyl)pyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (**8e**)




Yellow solid (66% yield), Chemical Formula: $C_{20}H_{19}BrN_4O_2$, 1H NMR (399 MHz, $cdCl_3$) δ 8.72 (d, $J = 7.9$ Hz, 1H), 8.01 (s, 1H), 7.69 (s, 1H), 7.25 (d, $J = 2.0$ Hz, 4H), 4.74 (d, $J = 24.6$ Hz, 2H), 3.64 (s, 2H), 3.34 (s, 4H), 2.95 (d, $J = 1.9$ Hz, 4H). ^{13}C NMR (100 MHz, $cdCl_3$) δ 167.2, 147.2, 143.5, 130.3, 126.7, 121.9, 120.1, 119.9, 117.4, 111.6, 109.2, 63.2, 45.38, 44.7, 35.2, 28.7.

1-Bromo-6-isopropylpyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (**8f**)




Off-white solid (76% yield), Chemical Formula: $C_{17}H_{14}BrN_3O$. 1H NMR (399 MHz,) δ 8.52 – 8.46 (m, 1H), 8.23 (d, $J = 7.2$ Hz, 1H), 8.03 (d, $J = 8.4$ Hz, 1H), 7.80 (d, $J = 9.2$ Hz, 1H), 7.36 (t, $J = 7.9$ Hz, 1H), 7.25 (d, $J = 1.2$ Hz, 1H), 6.88 (t, $J = 7.0$ Hz, 1H), 5.09 – 4.75 (m, 1H), 1.89 – 1.83 (m, 6H). ^{13}C NMR (100 MHz, $cdCl_3$) δ 143.1, 129.3, 128.1, 126.0, 122.1, 118.5, 116.0, 113.5, 49.6, 20.8, 19.8. HRMS (ESI) calcd for $C_{17}H_{14}BrN_3O$ m/z : 355.0320, found 355.0624.

6-Benzyl-1-bromopyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (**8g**)




Light yellow solid (82% yield), Chemical Formula: $C_{21}H_{14}BrN_3O$. 1H NMR (400 MHz, $cdCl_3$) δ 7.53 (dt, $J = 9.1, 1.2$ Hz, 1H), 7.42 (dd, $J = 3.7, 1.1$ Hz, 1H), 7.36 (dd, $J = 5.1, 1.1$ Hz, 1H), 7.21 – 7.04 (m, 6H), 7.01 (dt, $J = 6.8, 1.2$ Hz, 1H), 6.55 – 6.44 (m, 2H), 4.61 (t, $J = 7.9$ Hz, 2H). ^{13}C NMR (100 MHz, $cdCl_3$) δ 139.5, 135.5, 129.5, 129.5, 128.0, 127.4, 125.4, 123.6, 123.0, 119.6, 112.9, 47.4. HRMS (ESI) calcd for $C_{21}H_{14}BrN_3O$ m/z : 403.0320, found 403.0341.

1-Bromo-6-butylpyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (**8h**)




Yellow solid (84% yield), Chemical Formula: $C_{18}H_{16}BrN_3O$. 1H NMR (399 MHz, $cdCl_3$) δ 8.57 (dd, $J = 8.0, 1.4$ Hz, 1H), 8.36 (d, $J = 7.3$ Hz, 1H), 8.09 – 8.02 (m, 1H), 7.80 (d, $J = 9.3$ Hz, 1H), 7.42 – 7.33 (m, 1H), 7.25 (d, $J = 1.2$ Hz, 1H), 7.22 – 7.13 (m, 1H), 6.91 – 6.83 (m, 1H), 4.67 (t, $J = 7.8$ Hz, 2H), 1.92 (t, $J = 8.0$ Hz, 2H), 1.56 (q, $J = 7.5$ Hz, 2H), 1.04 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (100 MHz, $cdCl_3$) δ 160.6, 141.8, 139.1, 131.1, 130.3, 129.2, 127.2, 126.6, 123.3, 122.6, 120.1, 117.1, 113.3, 43.3, 31.9, 20.0, 13.8. HRMS (ESI) calcd for $C_{18}H_{16}BrN_3O$ m/z : 369.0477, found 369.0480.

1-Bromo-6-(2-morpholinoethyl)pyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (8i)




Deep yellow solid (76% yield), Chemical Formula: $C_{20}H_{19}BrN_4O_2$. 1H NMR (399 MHz, $cdCl_3$) δ 8.57 (dd, $J = 8.0, 1.4$ Hz, 1H), 8.36 (d, $J = 7.3$ Hz, 1H), 8.09 – 8.02 (m, 1H), 7.80 (d, $J = 9.3$ Hz, 1H), 7.42 – 7.33 (m, 1H), 7.22 – 7.13 (m, 1H), 6.91 – 6.83 (m, 1H), 4.45 (ddd, $J = 14.0, 7.3, 4.5$ Hz, 2H), 3.61 – 3.50 (m, 3H), 3.45 (d, $J = 7.1$ Hz, 4H), 2.62 – 2.52 (m, 3H), 2.29 (ddd, $J = 12.7, 7.1, 4.1$ Hz, 3H). ^{13}C NMR (125 MHz, Common NMR Solvents) δ 149.0, 132.7, 123.3, 118.2, 118.0, 117.0, 116.3, 115.9, 115.5, 115.3, 114.1, 107.0, 111.1, 100.3, 54.6, 43.3, 41.8, 35.3.

6-Butyl-9-chloropyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (8l)




White solid (68% yield), Chemical Formula: $C_{18}H_{16}ClN_3O$. 1H NMR (399 MHz, $cdCl_3$) δ 8.52 (dd, $J = 14.1, 8.1$ Hz, 2H), 8.32 (d, $J = 7.1$ Hz, 1H), 7.86 – 7.77 (m, 1H), 7.58 (t, $J = 7.5$ Hz, 1H), 7.32 – 7.22 (m, 1H), 7.25 (s, 2H), 7.25 (s, 2H), 6.82 (t, $J = 7.2$ Hz, 1H), 4.65 (t, $J = 7.9$ Hz, 2H), 1.95 – 1.86 (m, 2H), 1.64 – 1.55 (m, 2H), 1.04 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (100 MHz, $cdCl_3$) δ 133.0, 129.2, 127.5, 124.8, 122.5, 121.5, 113.0, 43.1, 32.2, 20.0, 13.8. HRMS (ESI) calcd for $C_{18}H_{16}ClN_3O$: 325.0982, found 325.0915.

9-Bromo-6-butylpyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (8m)




White solid (80% yield), Chemical Formula: $C_{18}H_{16}BrN_3O$. 1H NMR (399 MHz, $cdCl_3$) δ 8.51 (d, $J = 8.7$ Hz, 2H), 8.39 (d, $J = 7.9$ Hz, 1H), 7.81 (t, $J = 7.7$ Hz, 1H), 7.64 – 7.50 (m, 2H), 7.23 (dd, $J = 13.4, 3.7$ Hz, 2H), 4.63 (t, $J = 7.8$ Hz, 2H), 1.91 (q, $J = 7.8$ Hz, 2H), 1.12 – 1.03 (m, 3H). ^{13}C NMR (100 MHz, $cdCl_3$) δ 133.1, 129.4, 127.5, 126.7, 122.6, 121.9, 119.5, 42.6, 29.7, 19.9, 13.8. HRMS (ESI) calcd for $C_{18}H_{16}BrN_3O$ m/z : 369.0477, found 369.0480.

10-Bromo-6-butylpyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (8n)




White solid (84% yield), Chemical Formula: $C_{18}H_{16}BrN_3O$. 1H NMR (399 MHz, $cdCl_3$) δ 8.55 – 8.48 (m, 1H), 8.45 – 8.32 (m, 2H), 7.85 – 7.76 (m, 1H), 7.75 – 7.68 (m, 1H), 7.61 – 7.52 (m, 1H), 7.23 – 7.14 (m, 1H), 4.67 (t, $J = 7.9$ Hz, 2H), 1.92 (t, $J = 7.7$ Hz, 2H), 1.58 (h, $J = 7.4$ Hz, 2H), 1.16 (s, 2H), 1.08 – 1.00 (m, 3H). ^{13}C NMR (100 MHz, $cdCl_3$) δ 167.2, 147.2, 143.5, 130.3, 129.4, 126.7, 122.4, 121.9, 120.1, 119.9, 117.4, 111.6, 109.2, 48.2, 30.4, 20.2, 13.7. HRMS (ESI) calcd for $C_{18}H_{16}BrN_3O$ m/z : 369.0477, found 369.0480.

6-Isopropylpyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (8p)




White solid (72% yield), Chemical Formula: $C_{17}H_{15}N_3O$. 1H NMR (399 MHz, $cdCl_3$) δ 8.51 (d, 1H), 8.40 (d, 1H), 8.37 (d, 1H), 7.84 – 7.81 (m, 1H), 7.74 – 7.72 (m, 1H), 7.57 – 7.54 (m, 1H), 7.19 (t, 1H), 6.86 (t, $J = 7.7$ Hz, 1H), 4.93 (m, 1H), 1.87 (d, 6H). ^{13}C NMR (100 MHz, $cdCl_3$) δ 143.1, 136.3, 130.21, 128.5, 128.4, 126.1, 122.4, 117.6, 115.4, 112.1, 51.2, 24.2. HRMS (ESI) calcd for $C_{17}H_{15}N_3O$. m/z : 277.1215, found 277.1205.

6-Benzylpyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (8q)




Light yellow solid (78% yield), Chemical Formula: $C_{21}H_{15}N_3O$. 1H NMR (399 MHz, $cdCl_3$) δ 8.57 (d, $J = 8.0$ Hz, 1H), 8.48 (dd, $J = 16.3, 8.0$ Hz, 1H), 8.37 (d, $J = 8.0$ Hz, 1H), 8.26 (d, $J = 7.8$ Hz, 1H), 8.19 – 8.12 (m, 1H), 7.90 – 7.77 (m, 1H), 7.72 – 7.60 (m, 1H), 7.62 – 7.53 (m, 1H), 7.40 – 7.22 (m, 3H), 7.12 – 7.04 (m, 1H), 6.89 – 6.82 (m, 1H), 6.60 (t, $J = 7.0$ Hz, 1H), 4.62 (d, $J = 7.9$ Hz, 1H). ^{13}C NMR (100 MHz, $cdCl_3$) δ 133.4, 133.1, 129.6, 129.4, 129.4, 127.9, 127.4, 127.3, 125.5, 123.2, 122.9, 121.9, 117.6, 114.6, 46.9. HRMS (ESI) calcd for $C_{21}H_{15}N_3O$ m/z : 325.1215, found 325.1232.

6-Butylpyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (8r)




Deep yellow solid (80% yield), Chemical Formula: $C_{18}H_{17}N_3O$. 1H NMR (399 MHz, $cdCl_3$) δ 8.55 – 8.48 (m, 1H), 8.45 – 8.32 (m, 2H), 7.85 – 7.76 (m, 1H), 7.75 – 7.68 (m, 1H), 7.61 – 7.52 (m, 1H), 7.28 – 7.14 (m, 1H), 6.92 – 6.83 (m, 1H), 4.67 (t, $J = 7.9$ Hz, 2H), 1.92 (t, $J = 7.7$ Hz, 2H), 1.58 (h, $J = 7.4$ Hz, 2H), 1.08 – 1.00 (m, 3H). ^{13}C NMR (100 MHz, $cdCl_3$) δ 161.4, 143.0, 133.0, 131.6, 129.3, 127.2, 124.1, 123.5, 122.9, 122.8, 121.9, 119.2, 112.9, 42.8, 32.2, 20.0, 13.8. HRMS (ESI) calcd for $C_{18}H_{17}N_3O$ m/z : 291.1372, found 291.1369.

6-(2-Morpholinoethyl)pyrido[2',1':2,3]imidazo[4,5-c]isoquinolin-5(6H)-one (8s)






Yellow solid (60% yield), Chemical Formula: $C_{20}H_{20}N_4O_2$. 1H NMR (399 MHz, $cdCl_3$) δ 8.77 (s, 1H), 8.50 (d, $J = 8.2$ Hz, 1H), 8.42 (d, $J = 8.0$ Hz, 1H), 7.82 (t, $J = 7.5$ Hz, 1H), 7.72 (d, $J = 9.4$ Hz, 1H), 7.57 (t, $J = 7.7$ Hz, 1H), 7.25 (s, 2H), 6.89 (t, $J = 6.7$ Hz, 1H), 4.82 (s, 2H), 3.72 (s, 5H), 3.63 (s, 2H), 2.95 (s, 2H), 2.88 (s, 3H). ^{13}C NMR (100 MHz, $cdCl_3$) δ 168.1, 149.3, 143.2, 132.7, 131.6, 127.0, 126.4, 123.6, 117.1, 111.8, 111.2, 108.3, 66.6, 56.5, 52.1, 44.2.

^1H NMR, ^{13}C NMR spectra of products **6a** and **8**




^1H NMR of **6a**







^1H NMR of **8b**




^{13}C NMR of **8b**




¹H NMR of 8c




¹³C NMR of 8c




¹H NMR of 8d




¹³C NMR of 8d




^1H NMR of **8e**





^{13}C NMR of **8e**




¹H NMR of 8f




¹³C NMR of 8f







¹H NMR of 143




¹³C NMR of 8h




¹H NMR of 8i




¹³C NMR of 8i




¹H NMR of **8I**




¹³C NMR of **8I**




¹H NMR of **8m**





¹³C NMR of **8m**






¹H NMR of 8p




¹³C NMR of 8p







¹H NMR of 8s



¹³C NMR of 8s