

Supporting Information

for

Chiral multifunctional thiourea-phosphine catalyzed asymmetric [3 + 2] annulation of Morita–Baylis–Hillman carbonates with maleimides

Hong-Ping Deng¹, De Wang², Yin Wei¹ and Min Shi*^{1,2,§}


Address: ¹State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Road, Shanghai 200032, People's Republic of China and ²Key Laboratory for Advanced Materials and Institute of Fine Chemicals, School of Chemistry & Molecular Engineering, East China University of Science and Technology, and 130 MeiLong Road, Shanghai 200237, People's Republic of China

Email: Min Shi - Mshi@mail.sioc.ac.cn

§Fax: 86-21-64166128

*Corresponding author

Crystal structure data of compound 3r



The crystal data of **3r** have been deposited in CCDC with number 863228. Empirical Formula: $C_{16}H_{14}BrNO_3$; Formula Weight: 348.19; Crystal Color, Habit: colorless, Crystal Dimensions: $0.211 \times 0.157 \times 0.104$ mm; Crystal System: Orthorhombic; Lattice Parameters: $a = 9.5280(6)\text{\AA}$, $b = 9.6271(6)$ \AA , $c = 16.3779(10)$ \AA , $\alpha = 90^\circ$, $\beta = 90^\circ$, $\gamma = 90^\circ$, $V = 1502.30(16)$ \AA^3 ; Space group: $P2(1)2(1)2(1)$; $Z = 4$; $D_{calc} = 1.539$ g/cm^3 ; $F_{000} = 704$; Final R indices [$I > 2\sigma(I)$]: $R_1 = 0.0384$; $wR_2 = 0.0900$.