

An exceptional P-H phosphonite: Biphenyl-2,2'-bisfenchylchlorophosphite and derived ligands (BIFOPs) in enantioselective copper-catalyzed 1,4-additions

T. Kop-Weiershausen, J. Lex, J.-M. Neudörfl and B. Goldfuss*

Full Research Paper

Open Access

Address:
Institut für Organische Chemie, Universität zu Köln, Greinstrasse 4,
50939 Köln, Germany

Email:
T. Kop-Weiershausen - t.kop-weiershausen@uni-koeln.de; J. Lex -
johann.lex@uni-koeln.de; J.-M. Neudörfl - aco48@uni-koeln.de;
B. Goldfuss* - Goldfuss@Uni-Koeln.de

* Corresponding author

Keywords:
phosphorus ligands; chirality; biaryls; asymmetric conjugate additions;
phosphoramidites; phosphites; phosphonites; X-ray analyses

Beilstein Journal of Organic Chemistry **2005**, 1, No. 6.
doi:10.1186/1860-5397-1-6

Received: 29 June 2005
Accepted: 26 August 2005
Published: 26 August 2005

© 2005 Kop-Weiershausen et al; licensee Beilstein-Institut.
License and terms: see end of document.




Abstract

Biphenyl-2,2'-bisfenchol (BIFOL) based chlorophosphite, BIFOP-Cl, exhibits surprisingly high stabilities against hydrolysis as well as hydridic and organometallic nucleophiles. Chloride substitution in BIFOP-Cl proceeds only under drastic conditions. New enantiopure, sterically demanding phosphorus ligands such as a phosphoramidite, a phosphite and a P-H phosphonite (BIFOP-H) are hereby accessible. In enantioselective Cu-catalyzed 1,4-additions of ZnEt₂ to 2-cyclohexen-1-one, this P-H phosphonite (yielding 65% ee) exceeds even the corresponding phosphite and phosphoramidite.

Introduction

Chiral monodentate phosphorus ligands with C₂-symmetric diol backbones, e.g. with the prominent BINOLs or TADDOLs, are fundamental for the construction of efficient enantioselective transition metal catalysts, especially for copper-catalyzed 1,4-additions. [1-18] Such asymmetric conjugate additions of diethylzinc to enones are often highly enantioselective, espe-

cially with phosphoramidites (amidophosphites) and phosphites. [2,19-41] These chiral ligands (L*) exhibit large steric demands and good metal to ligand back bonding abilities. Such ligands generate active R-Cu^I-L* catalysts and support the rate determining reductive elimination in the catalytic cycle (Scheme 1). [42-47]




Common basis for diol-based phosphoramidites and phosphites are highly reactive chlorophosphites. [48-51] These intermediates are converted usually *in situ* with amines or alcohols to the modular, enantiopure ligands (Scheme 2). [2,19-41].

Modular fencholates were recently applied in chiral organolithium reagents [52-58] and in organozinc [59-62] as well as in organopalladium catalysts [63-67] to study origins of enantioselectivities in C-C-couplings. The rigid, terpene-based bicyclo[2.2.1]heptane unit enables efficient, stereoselective access to crystalline diol ligands such as BIFOL (*biphenylbisfenchol*), [68-72] which we here apply for constructions of

new BIFOL-based, phosphorus ligands, i.e. *biphenylbisfenchol*phosphanes (BIFOPs). BIFOPs with high steric demand and good acceptor abilities are then employed in enantioselective Cu-catalyzed 1,4-additions.

Results and Discussion

Coupling of bis-lithiated biphenyl-2,2'-bisfenchol (BIFOL), synthesized from 2,2'-dilithiobiphenyl and (-)-fenchone, [68-72] with PCl₃ or PBr₃ yields the enantiopure halophosphites BIFOP-Cl, **1** (62% yield) and BIFOP-Br, **2** (69% yield, Scheme 3), which are air stable (no hydrolysis or oxidation) over weeks, crystalline and analyzable via X-ray diffraction. [73-76] In



Scheme 4: Geometries of BIFOP-systems with respect to biaryl dihedral angles ($C_2-C_1-C_1'-C_2'$, BAA), fenchyl-aryl dihedral angles ($C_1-C_2-C_3-O_1$) on the lone pair-side of phosphorus (FAA-lp) and at the substituent (X) side (FAA), the pyramidalicity of phosphorus measured as angle sum and the distance of phosphorus to the biaryl axis (C_1-C_1').

close analogy to the hydrogen bonded *M*-BIFOL, [68-72] only *minus*-(*M*)-conformations of biaryl axes are found in these BIFOP (*biphenylbisfencholphosphane*) halides (Figure 1 and Figure 2).

Surprisingly, the halophosphites **1** (BIFOP-Cl) and **2** (BIFOP-Br) are very reluctant in reactions with various nucleophilic reagents to give halide substitution (Table 1). [77,78] No nucleophilic substitution [2,19-41] is observed for **1** in treatments with equimolar suspensions of $LiAlH_4$ in hexanes or THF at 25°C for 3 h (Table 1, entries 1 and 2). Only elevated temperat-

ures (69°C), longer reaction times (12 h) and an excess of $LiAlH_4$ yields the P-H phosphonite BIFOP-H, **3** (79% yield, Table 1, entry 3, Figure 3). Even the strong C-nucleophilic reagents methylolithium, ethyllithium, *n*-butyllithium and *t*-butyllithium gave no expected conversions at low temperatures. Ethylation of BIFOP-Cl (**1**) yielding BIFOP-Et (**4**) was observed with ethyllithium only at elevated temperatures (Table 1, entry 7) or with a large excess of diethylzinc at room temperature (Table 1, entry 13). Similarly, **1** was converted to BIFOP-*n*Bu (**5**, Figure 4) only with an excess of *n*-BuLi at elevated temperatures. The resistance of BIFOP-Cl (**1**) to O- and

Table 1: Reactivity of BIFOP-Cl (**1**) with various nucleophilic reagents.

entry	Reagent	1 : reagent	conditions	yield ^a
1	$LiAlH_4$	1 : 1	Rt, 3 h, THF	- ^b
2	$LiAlH_4$	1 : 1	Rt, 3 h, hexanes	- ^b
3	$LiAlH_4$	1 : 2.5	reflux, 12 h, hexanes	3 (79%)
4	MeLi	1 : 1.2 ^c	-78°C, hexanes, Et_2O^d	- ^b
5	MeLi	1 : 5 ^c	reflux, 24 h hexanes, Et_2O	- ^b
6	EtLi	1 : 1.2 ^c	-78°C, hexanes, benzene ^d	- ^b
7	EtLi	1 : 5 ^c	reflux, 24 h, hexanes, benzene	4 (63%)
8	<i>n</i> BuLi	1 : 1.2	-78°C-rt, hexanes ^d	- ^b
9	<i>n</i> BuLi	1 : 5	reflux, 24 h, hexanes	5 (74%)
10	<i>t</i> BuLi	1 : 1.2	-78°C-rt, hexanes ^d	- ^b
11	<i>t</i> BuLi	1 : 5	reflux, 48 h, hexanes	- ^b
12	$ZnEt_2$	1 : 2	rt, 2 h, toluene	- ^b
13	$ZnEt_2$	1 : 140	rt, 2 h, toluene	4 (89%)
14	H_2O	1 : 110	rt, 3 h	- ^b
15	H_2O/KOH	1 : 220 ^e	reflux, 5 days	8 (98%)
16	LiOPh	1 : 5	-78°C-rt, hexanes ^d	- ^b
17	LiOPh	1 : 5	reflux, 24 h, hexanes	6 (47%)
18	$LiNEt_2$	1 : 5	-78°C-rt, hexanes ^d	- ^b
19	$LiNEt_2$	1 : 5	reflux, 24 h, hexanes	7 (47%)

^a The reaction of **1** ($\delta^{31}P$: 154.4) was monitored by ^{31}P -NMR-spectroscopy; isolated yields are given;

^b Only pure BIFOP-Cl (**1**) was recovered (> 91%);

^c MeLi 1.6 M solution in diethylether; EtLi 0.5 M solution in benzene/cyclohexane (90/10);

^d Reactions were performed at -78°C for 3 h and subsequently at rt for 3 h;

^e With KOH (1 M, 0.01 g, 0.18 mmol) in water.




Figure 1: X-ray crystal structure of **BIFOP-Cl (1)**. Distances are given in Å. (BAA = biaryl angle between $C_2-C_1-C_1'-C_2'$; FAA-lp = fenchyl-aryl dihedral angle between $C_1-C_2-C_3-O_1$; FAA = fenchyl-aryl dihedral angle between $C_1'-C_2'-C_3'-O_2$). The probability of ellipsoids is 40% (CDC 270531).




Figure 2: X-ray structures of **BIFOP-Br (2)**. Distances are given in Å. (BAA = biaryl angle between $C_2-C_1-C_1'-C_2'$; FAA-lp = fenchyl-aryl dihedral angle between $C_1-C_2-C_3-O_1$; FAA = fenchyl-aryl dihedral angle between $C_1'-C_2'-C_3'-O_2$). The probability of ellipsoids is 30% (CCDC 270532).

N-nucleophiles is apparent from reactions with H_2O , $LiOPh$ and $LiNEt_2$. While no hydrolysis of **1** is observed at ambient temperature, only reflux and basic conditions (KOH) yield complete hydrolysis of **1** to **BIFOP(O)-H (8)** (98%, Table 1, entry 14, Figure 5). [79,80] The phosphite **BIFOP-OPh (6)** (40%, Figure 6) and the phosphoramidite **BIFOP-NEt₂ (7)** (47%, Figure 7) are accessible from **1** only at elevated temperatures with $LiOPh$ or $LiNEt_2$. The oxo-derivative **BIFOP(O)-Cl (9)** is synthesized by coupling of **BIFOL** with $POCl_3$ (65%, Figure 8).

The high steric demand of the embedding fenchane units provides explanations for the unexpectedly low reactivity of the $>P-Cl$ moiety in **BIFOP-Cl (1)**. The geometries of all **BIFOP**-derivatives are remarkable with respect to their biaryl-angles, the fenchyl-aryl-angles, the pyramidity at the phosphorus atoms as well as the positions of the phosphorus atom in the hydrophobic fenchane cavities (Scheme 4, Table 2).

A strong preference for the *minus (M)*-biaryl conformation was found in **BIFOL** (Scheme 3) and was attributed to hydrogen bond linked chiral fenchane units, in the solid state and in solution. [68-72] Similarly, all phosphorus linked **BIFOPs** exhibit *M*-biaryl axes with dihedral angles varying from -91° to -99° (Scheme 4, Table 2). The alternative *plus (P)*-conformations were not found experimentally, they are computed to be disfavored by ca. 20 to 35 kcal mol⁻¹ (Table 2). As in *M*-**BIFOL**, [68-72] the strong destabilization of these *plus-(P)*-conformations arise from steric repulsion of *endo*-oriented fenchane units in **BIFOPs**, *endo*-methyl groups are close to the phosphorus atoms (Figure 9).

Free rotation around the fenchyl aryl bonds is hindered in aryl fenchol derivatives by the methyl-methylene-bifurcation of the fenchane scaffolds (Scheme 4). These fenchyl aryl dihedral angles ($O_1-C_2-C_3-C_1$, FAA, Scheme 4) are crucial for the "bite" of the chiral diol unit and are constrained between 30° and 46° (Table 2), similar to fenchyl aryl angles previously analyzed in lithium fencholates. [54,55] In **BIFOL**, the asymmetry of the H-bond gives rise to two different fenchyl aryl angles (25° and 31° , Table 2). [68-72] Likewise, the pyramidity at the phosphorus atoms in **BIFOPs** distorts the inherent C_2 -symmetry of the biphenylbisfenchol units to asymmetry (C_1). Fenchanes close to the phosphorus lone pairs (with FAA-lp) can be differentiated from fenchanes close to substituents at phosphorus (with FAA, Scheme 4). The difference between these two FAA dihedral angles is a measure for the **BIFOP**-asymmetry, which is small for **BIFOP-H (3)** and **BIFOP-NEt₂ (7)**, but large for **BIFOP-Cl (1)** and **BIFOP-OPh (6)**, Table 2).

The phosphorus atoms, essential for coordination to (late transition) metals, exhibit slightly different degrees of pyramidity,




Figure 3: X-ray structures of BIFOP-H (**3**). Distances are given in Å. (BAA = biaryl angle between $C_2-C_1-C_1'-C_2'$; FAA-lp = fenchyl-aryl dihedral angle between $C_1-C_2-C_3-O_1$; FAA = fenchyl-aryl dihedral angle between $C_1'-C_2'-C_3'-O_2$). The probability of ellipsoids is 40% (CCDC 270533).




Figure 5: X-ray structures of BIFOP(O)-H (**8**). Distances are given in Å. (BAA = biaryl angle between $C_2-C_1-C_1'-C_2'$; FAA-lp = fenchyl-aryl dihedral angle between $C_1-C_2-C_3-O_1$; FAA = fenchyl-aryl dihedral angle between $C_1'-C_2'-C_3'-O_2$). The probability of ellipsoids is 40% (CCDC 270537).




Figure 4: X-ray structures of BIFOP-*n*Bu (**5**). Distances are given in Å. (BAA = biaryl angle between $C_2-C_1-C_1'-C_2'$; FAA-lp = fenchyl-aryl dihedral angle between $C_1-C_2-C_3-O_1$; FAA = fenchyl-aryl dihedral angle between $C_1'-C_2'-C_3'-O_2$). The probability of ellipsoids is 40% (CCDC 270534).




Figure 6: X-ray structures of phosphite BIFOP-OPh (**6**). Distances are given in Å. (BAA = biaryl angle between $C_2-C_1-C_1'-C_2'$; FAA-lp = fenchyl-aryl dihedral angle between $C_1-C_2-C_3-O_1$; FAA = fenchyl-aryl dihedral angle between $C_1'-C_2'-C_3'-O_2$). The probability of ellipsoids is 40% (CCDC 270535).

as it is measured by angle sums (planarity) from 300° to 309° (Table 2, Scheme 4). The degree of encapsulation of the phosphorus atoms by the fenchane units is measured by the distance (d) of the phosphorus atoms to the center of the biaryl axes (C_1-


C_1' , Scheme 4). The tightest encapsulation and fenchane embedding of phosphorus atoms is apparent for the halophosphites BIFOP-Cl (**1**, 2.471 Å) and BIFOP-Br (**2**, 2.476 Å, Table 2), explaining their unusual low reactivity (Table 1).

Table 2: X-ray structure geometries (cf. Scheme 4) and computed energies of BIFOL and BIFOP-X (1–7).

BIFOP (-X)	E _{rel} (kcal mol ⁻¹) ^a	BAA (°) ^b	angle sum (°) ^c	FAA-lp (°)	FAA (°)	d (Å) ^d
M-BIFOL	+12.5	-95.0 ^e	-	(25)	(31)	-
1 (-Cl)	+28.4	-91.3	305.2	37.9	46.3	2.471
2 (-Br)	+35.0	-91.7	306.6	38.9	36.9	2.476
3 (-H)	+27.9	-98.2	301.4	30.0	29.7	2.669
5 (-nBu)	+26.3	-99.4	303.6	31.4	34.5	2.792
6 (-OPh)	+27.7	-93.3	300.0	34.5	44.8	2.567
7 (-NEt ₂)	+22.5	-91.1	309.2	35.8	35.1	2.877

^a Relative Destabilization of *plus* (*P*) conformations according to B3LYP/6-31G*//PM3 computations;^b Biaryl dihedral angle between C2-C1-C1'-C2' atoms (BAA) in degree;^c Angle sum at phosphorus atom (pyramidal) in degree;^d Distance (d) between phosphorus atoms and the center of the biaryl axis (C1-C1');^e Hydrogend bonded *M*-conformer.**Table 3:** Enantioselective Cu-catalyzed 1,4-additions of diethylzinc to 2-cyclohexen-1-one (cf. Scheme 5).^a

L*, i.e. BIFOP(O)-X	substituent X	yield (%) ^b	%ee (config.) ^c
1 ^d	Cl	98 ^d	11 (<i>R</i>) ^d
3	H	92	65 (<i>R</i>)
4	Et	98	11 (<i>R</i>)
5	<i>n</i> Bu	97	7 (<i>R</i>)
6	OPh	73	16 (<i>R</i>)
7	NEt ₂	98	12 (<i>R</i>)
8	(O)-H	89	2 (<i>R</i>)
9	(O)-Cl	87	3 (<i>S</i>)

^a Reaction conditions: -20°C, 3 h in CH₂Cl₂; L*:Cu(OTf)₂ ratio (2:1);^b Yield determined by GC;^c The ee's are determined by GC with the chiral column lipodex E 0.2 μm, 50 m, 0.25 mm;^d Chlorophosphite **1** was converted to P-Et phosphonite BIFOP-Et, **4**.**Scheme 5:** Biphenyl-2,2'-bisfenchol based phosphanes (BIFOPs) as chiral ligands in enantioselective Cu-catalyzed 1,4-additions.




Figure 7: X-ray structures of phosphoramidite **BIFOP-NEt₂** (**7**). Distances are given in Å. (BAA = biaryl angle between C₂-C₁-C₁'-C₁'; FAA-lp = fenchyl-aryl dihedral angle between C₁-C₂-C₃-O₁; FAA = fenchyl-aryl dihedral angle between C₁'-C₂'-C₃'-O₂). The probability of ellipsoids is 40% (CCDC 270536).




Figure 8: X-ray structures of **BIFOP(O)-Cl** (**9**). Distances are given in Å. (BAA = biaryl angle between C₂-C₁-C₁'-C₂'; FAA-lp = fenchyl-aryl dihedral angle between C₁-C₂-C₃-O₁; FAA = fenchyl-aryl dihedral angle between C₁'-C₂'-C₃'-O₂). The probability of ellipsoids is 40% (CCDC 270538).



Cu-catalyzed, enantioselective 1,4-additions of diethylzinc to 2-cyclohexene-1-one were employed as test reactions for the monodentate phosphorus ligands BIFOPs **1** and **3–7** as well as the oxo-derivatives **8** and **9** (Table 3, Scheme 5).

The low reactivity of the chlorophosphite **1** with metal organic nucleophiles (Table 1) points to its potential suitability as ligand for late, electron-rich transition metals, such as Cu^I. The rate determining reductive elimination was expected to be favored by the good metal to ligand back bonding properties of the $\sigma^*(\text{P}-\text{Cl})$ acceptor as is well established in phosphites, i.e. $\sigma^*(\text{P}-\text{OR})$, and phosphoramidites, i.e. $\sigma^*(\text{P}-\text{NR}_2)$. Computed anharmonic CO-frequencies on Cu^I-model complexes indeed point to highest ν_{CO} stretching frequencies for P-ligands with strongest acceptor character, i.e. the halophosphites (Scheme 6). [81–85]

However, under catalysis conditions, BIFOP-Cl (**1**) converts to the ethylphosphonite BIFOP-Et (**4**), which yields *R*-3-ethylcyclohexanone with 11% ee (Table 3). Apparently, the higher Lewis acidity of organozincs supports faster nucleophilic substitution in (**1**) than with organolithiums (Table 1). Unprecedented however is the P-H phosphonite BIFOP-H (**3**), which yields with 65 % ee a much higher enantioselectivity than the corresponding phosphite (BIFOL-OPh, 16%ee) and phosphoramidite (BIFOL-NEt₂, 12%ee, Table 3). A good back bonding characteristic between halophosphites and phosphites is indeed apparent for the P-H unit (Scheme 6). [86,87] The relative high enantioselectivity of **3** is remarkable, as the asymmetry of **3**, measured from the difference of its fenchyl-aryl angles, is rather small (FAA-lp = 30.0° vs. FAA = 29.7°, Table 2). The phosphorus atom in **3** is only slightly encapsulated by the fenchane moieties, due to the rather long *d*-distance (Scheme 4, Table 2, 2.7 Å) and the low steric shielding by the H-atom. Indeed, P-H phosphonite **3** coordinates tightly to Cu^{II}-ions. Metal free BIFOP-H (**3**) gives in CDCl₃ a ³¹P-NMR signal at $\delta = 139.6$ with a ¹J (P,H) coupling of 214.5 Hz. With half of an equivalent of Cu(OTf)₂, no free **1** is detectable, only a [(**1**)₂Cu(OTf)₂] complex is evident from a ³¹P-NMR signal at $\delta = 81.1$ with a stronger ¹J (P,H) coupling of 299.5 Hz. [86,89]

Conclusion

The large steric demand of embedding fenchane units makes phosphorus atoms in BIFOPs hardly accessible by nucleophilic reagents and leads to an unusually high stability, e.g. for the chlorophosphite BIFOP-Cl (**1**). While **1** converts to the P-Et phosphonite BIFOP-Et (**4**) during Cu-catalyzed 1,4-additions of diethylzinc to cyclohexenone, the P-H phosphonite BIFOP-H (**3**) is stable and gives even a higher enantioselectivity than a corresponding phosphite or phosphoramidite. Hence, the large steric demand and the relatively low accessibility of the phosphorus atoms in biphenyl-2,2'-bisfenchylphosphites (BIFOPs)



finds the special suitability for BIFOP-H (**3**) as P-H phosphonite ligand in transition metal catalysis. This points to promising applications of **3** or analogue P-H ligands in enantioselective catalysis.

Supporting Information

Supporting Information File 1

Experimental details

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-1-6-S1.doc>]

Acknowledgments

We are grateful to the Fonds der Chemischen Industrie for financial support as well as for a Dozenten-Stipendium to B. G. We also thank the Deutsche Forschungsgemeinschaft (DFG) for support (GO-930/9, GO-930/7 and GO-930/5). We are grateful to the Bayer AG, the BASF AG, the Wacker AG, the Degussa AG, the Raschig GmbH, the Symrise GmbH, the Solvay GmbH and the OMG AG for generous gifts of laboratory equipment and chemicals.

References

- d'Augustin, M.; Palais, L.; Alexakis, A. *Angew. Chem.* **2005**, *117*, 1440. *Angew. Chem., Int. Ed.* **2005**, *44*, 1376.
- Sebesta, R.; Pizzuti, M. G.; Boersma, A. J.; Minnaard, A. J.; Feringa, B. L. *Chem. Commun.* **2005**, 1711.
- Shi, M.; Zhang, W. *Adv. Synth. Catal.* **2005**, *347*, 535.
- Wu, X.-Y.; Li, X.-G.; Zhao, G. *Lett. Org. Chem.* **2005**, *2*, 65.
- Bonini, B. F.; Capito, E.; Comes-Franchini, M.; Ricci, A.; Bottoni, A.; Bernardi, F.; Miscione, G.; Giordano, P.; Cowley, L. A. R. *Eur. J. Org. Chem.* **2004**, *21*, 4442.
- Telfer, S. G.; Kuroda, R. *Coord. Chem. Rev.* **2003**, *242*, 33.
- Hayashi, T.; Yamasaki, K. *Chem. Rev.* **2003**, *103*, 2829.
- Alexakis, A.; Benhaim, C. *Eur. J. Org. Chem.* **2002**, *19*, 3221.
- Feringa, B. L.; Naasz, R.; Imbos, R.; Arnold, L. A. Copper-catalyzed Enantioselective Conjugate Addition Reactions of Organozinc Reagents. In *Modern Organocopper Chemistry*; Krause, N., Ed.; Wiley-VCH: Weinheim, 2002; pp 224 ff.
- Krause, N.; Hoffmann-Roder, A. *Synthesis* **2001**, *2*, 171.
- Krause, N. In *Organic Synthesis Highlights IV*; Schmalz, H.-G., Ed.; Wiley-VCH: Weinheim, 2000; pp 182 ff.
- Tomioka, K. Supplement to Chapter 31.1: Conjugate addition of organometallics to activated olefins. In *Asymmetric Catalysis, Supplement 2*; Jacobsen, E. N.; Pfaltz, A.; Yamamoto, H., Eds.; Springer: Berlin, Germany, 2004; pp 109–124.
- Yamanoi, Y.; Imamoto, T. *J. Org. Chem.* **1999**, *64*, 2988.
- Yan, M.; Yang, L.-W.; Wong, K.-Y.; Chan, A. S. C. *Chem. Commun.* **1999**, 11.
- Zhang, F.-Y.; Chan, A. S. C. *Tetrahedron: Asymmetry* **1998**, *9*, 1179.
- Krause, N. *Angew. Chem.* **1998**, *110*, 295. *Angew. Chem., Int. Ed.* **1998**, *37*, 283.
- Krause, N.; Gerold, A. *Angew. Chem.* **1997**, *109*, 194. *Angew. Chem., Int. Ed.* **1997**, *36*, 186.
- Knöbel, A. K. H.; Escher, I. H.; Pfaltz, A. *Synlett* **1997**, *36*, 2620. For Rh-catalyzed 1,4-additions.
- Hilgraf, R.; Pfaltz, A. *Adv. Synth. Catal.* **2005**, *347*, 61.
- Zeng, Q.-H.; Hu, X.-P.; Duan, Z.-C.; Liang, X.-M.; Zheng, Z. *Tetrahedron: Asymmetry* **2005**, *16*, 1233.

21. Reetz, M. T.; Ma, J.-A.; Goddard, R. *Angew. Chem.* **2005**, *117*, 416. *Angew. Chem., Int. Ed. Engl.* **2005**, *44*, 412.
22. Hu, X.-P.; Zheng, Z. *Org. Lett.* **2005**, *7*, 419.
23. Pianulli, U.; Daubos, P.; Clavarie, C.; Monti, C.; Gennai, C. *Eur. J. Org. Chem.* **2005**, 895.
24. Pfaltz, A. *Chimia* **2004**, *58*, 49.
25. Tissot-Croset, K.; Polet, D.; Alexakis, A. *Angew. Chem.* **2004**, *116*, 2480. *Angew. Chem., Int. Ed.* **2004**, *43*, 2426.
26. Elkema, R.; van Delden, R. A.; Feringa, B. L. *Angew. Chem.* **2004**, *116*, 5123. *Angew. Chem., Int. Ed. Engl.* **2004**, *43*, 5013.
27. Duursma, A.; Boiteau, J.-G.; Lefort, L.; Boogers, J. A. F.; de Vries, A. H. M.; de Vries, J. G.; Minnaard, A. J.; Feringa, B. L. *J. Org. Chem.* **2004**, *69*, 8045.
28. Alexakis, A.; Polet, D.; Rosse, S.; March, S. *J. Org. Chem.* **2004**, *69*, 5660.
29. Choi, H.; Hua, Z.; Ojima, I. *Org. Lett.* **2004**, *6*, 2689.
30. Alexakis, A.; Polet, D.; Rosset, S.; March, S. *J. Org. Chem.* **2004**, *69*, 5660.
31. Kinoshita, N.; Marx, K. H.; Tanak, K.; Tsubaki, K.; Kawabata, T.; Yoshikai, N.; Nakamura, E.; Fuji, K. *J. Org. Chem.* **2004**, *69*, 7960.
32. Hannen, P.; Militzer, H.-C.; Vogl, E. M.; Rampf, F. F. A. *Chem. Commun.* **2003**, 2210.
33. Pianulli, U.; Clavarie, C.; Daubos, P.; Gennai, C.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2003**, *5*, 4493.
34. Kočovský, P.; Vyskočil, Š.; Sorcina, M. *Chem. Rev.* **2003**, *103*, 3213.
35. Chen, Y.; Yekta, S.; Yudin, A. *Chem. Rev.* **2003**, *103*, 3155.
36. Ansell, J.; Wills, M. *Chem. Soc. Rev.* **2002**, *31*, 259.
37. Blume, F.; Zemolka, S.; Fey, T.; Kranich, R.; Schmalz, H.-G. *Adv. Synth. Catal.* **2002**, *344*, 868.
38. Seebach, D.; Beck, A. K.; Heckel, A. *Angew. Chem.* **2001**, *113*, 96. *Angew. Chem., Int. Ed.* **2001**, *40*, 92.
39. Naasz, R.; Arnold, L. A.; Minnaard, A. J.; Feringa, B. L. *Angew. Chem.* **2001**, *113*, 953. *Angew. Chem., Int. Ed.* **2001**, *40*, 927.
40. Alexakis, A.; Burton, J.; Vastra, J.; Benhaim, C.; Fournioux, X.; van den Heuvel, A.; Leveque, J.-M.; Maze, F.; Rosset, S. *Eur. J. Org. Chem.* **2000**, 4011.
41. Feringa, B. L. *Acc. Chem. Res.* **2000**, *33*, 346.
42. Nakanishi, W.; Yamanaka, M.; Nakamura, E. *J. Am. Chem. Soc.* **2005**, *127*, 1446.
43. Pfretzschner, T.; Kleemann, L.; Janza, B.; Harms, K.; Schrader, T. *Chem.–Eur. J.* **2004**, *10*, 6048.
44. Nakamura, E.; Mori, S. *Angew. Chem.* **2000**, *112*, 3902. *Angew. Chem., Int. Ed.* **2000**, *39*, 3750.
45. John, M.; Auel, C.; Behrens, C.; Marsch, M.; Harms, K.; Bosold, F.; Gschwind, R. M.; Rajamohanam, P. R.; Boche, G. *Chem.–Eur. J.* **2000**, *6*, 3060.
46. Mori, S.; Nakamura, E. *Chem.–Eur. J.* **1999**, *5*, 1534.
47. Canisius, J.; Gerold, A.; Krause, N. *Angew. Chem.* **1999**, *111*, 1727. *Angew. Chem., Int. Ed.* **1999**, *38*, 1644.
48. Alezra, V.; Bernardinelli, G.; Corminboeuf, C.; Frey, U.; Kündig, E. P.; Merbach, A. W.; Saudan, S. M.; Viton, F.; Weber, J. *J. Am. Chem. Soc.* **2004**, *126*, 4843.
49. van der Slot, S. C.; Duran, J.; Luten, J.; Kamer, P. C. J.; van Leeuwen, P. W. N. M. *Organometallics* **2002**, *21*, 3873.
50. Kündig, E. P.; Bourdin, B.; Bernardinelli, G. *Angew. Chem.* **1994**, *106*, 1931. *Angew. Chem., Int. Ed.* **1994**, *33*, 1856.
51. Burford, N.; Herbert, D. E.; Ragogna, P. J.; McDonald, R.; Ferguson, M. J. *J. Am. Chem. Soc.* **2004**, *126*, 17067.
52. Goldfuss, B. *Synthesis* **2005**, 2271.
53. Goldfuss, B.; Steigelmann, M.; Löschmann, T.; Schilling, G.; Rominger, F. *Chem.–Eur. J.* **2005**, *11*, 4019.
54. Goldfuss, B. Enantioselective addition of organolithiums to C=O groups. In *Organolithiums in Enantioselective Synthesis*; Hodgson, D. M., Ed.; Topics in Organometallic Chemistry, Vol. 5; Springer: Berlin, 2003; pp 21–35.
55. Goldfuss, B.; Steigelmann, M.; Rominger, F.; Urtel, H. *Chem.–Eur. J.* **2001**, 4456.
56. Goldfuss, B.; Steigelmann, M.; Rominger, F. *Angew. Chem.* **2000**, *112*, 4299. *Angew. Chem., Int. Ed.* **2000**, *39*, 4133.
57. Goldfuss, B.; Khan, S. I.; Houk, K. N. *Organometallics* **1999**, *18*, 2927.
58. Goldfuss, B.; Eisenträger, F. *Aust. J. Chem.* **2000**, *53*, 209.
59. Steigelmann, M.; Nisar, Y.; Rominger, F.; Goldfuss, B. *Chem.–Eur. J.* **2002**, *8*, 5211.
60. Goldfuss, B.; Steigelmann, M.; Khan, S. I.; Houk, K. N. *J. Org. Chem.* **2000**, *65*, 77.
61. Goldfuss, B.; Steigelmann, M. *J. Mol. Model.* **2000**, *6*, 166.
62. Goldfuss, B.; Steigelmann, M.; Rominger, F. *Eur. J. Org. Chem.* **2000**, 1785.
63. Goldfuss, B.; Löschmann, T.; Rominger, F. *Chem.–Eur. J.* **2004**, *10*, 5422.
64. Vázquez, J.; Goldfuss, B.; Helmchen, G. *J. Organomet. Chem.* **2002**, *641*, 67.
65. Kollmar, M.; Steinhagen, H.; Janssen, J. P.; Goldfuss, B.; Malinovskya, S. A.; Vázquez, J.; Rominger, F.; Helmchen, G. *Chem.–Eur. J.* **2002**, *8*, 3103.
66. Kollmar, M.; Goldfuss, B.; Reggelin, M.; Rominger, F.; Helmchen, G. *Chem.–Eur. J.* **2001**, *7*, 4913.
67. Goldfuss, B.; Kazmaier, U. *Tetrahedron* **2000**, *56*, 6493.
68. Goldfuss, B.; Rominger, F. *Tetrahedron* **2000**, *56*, 881.
69. Panev, S.; Linden, A.; Dimitrov, V. *Tetrahedron: Asymmetry* **2001**, *12*, 1313.
70. Dimitrov, V.; Linden, A.; Hesse, M. *Tetrahedron: Asymmetry* **2001**, *12*, 1331.
71. Dimitrov, V.; Dobrikov, G.; Genov, M. *Tetrahedron: Asymmetry* **2001**, *12*, 1323.
72. Dimitrov, V.; Rentsch, G. H.; Linden, A.; Hesse, M. *Helv. Chim. Acta* **2003**, *86*, 106.
73. Johes, P. G.; Kunze, C.; Thonnessen, H.; Schmutzler, R. *Acta Crystallogr., Sect. E* **2002**, *58*, 1432.
74. Chandrasekaran, A.; Day, R. O.; Sood, P.; Timosheva, N. V.; Sherlock, D. J.; Holmes, R. R. *Phosphorus, Sulfur Silicon Relat. Elem.* **2000**, *160*, 1.
75. Sherlock, D. J.; Chandrasekaran, A.; Day, R. O.; Holmes, R. R. *Inorg. Chem.* **1997**, *36*, 5082.
76. Cattani-Lorente, M.; Geoffroy, M.; Mishra, S. P.; Weber, J.; Bernardinelli, G. *J. Am. Chem. Soc.* **1986**, *108*, 7148. BIFOP-Cl (**1**) and BIFOP-Br (**2**) appear to be the first chiral halophosphites, which are characterized by X-ray-analysis.
77. Oh-Seuk, L.; Kiyull, Y.; Keum Duck, K.; In Sun, K.; Chan-Kyung, K.; Ikchoon, L. *J. Comput. Chem.* **2004**, *25*, 1740.
78. Haake, P.; Ossip, P. S. *J. Am. Chem. Soc.* **1971**, 6924.
79. Jiang, X.-b.; Minnaard, A. J.; Hessen, B.; Feringa, B. L.; Duchateau, A. L. L.; Andrien, J. G. O.; Boogers, J. A. F.; de Vries, J. G. *Org. Lett.* **2003**, *5*, 1503.

80. Lam, T. C. H.; Mak, W.-L.; Wong, V.-L.; Kwong, H.-L.; Sung, H. H. Y.; Lo, S. M. F.; Williams, I. D.; Leung, W.-H. *Organometallics* **2004**, *23*, 1247.
81. *Gaussian 03*, Revision C.02; Gaussian, Inc.: Wallingford, CT, 2004.
All computed structures were fully optimized and characterized by harmonic or anharmonic frequency computations using GAUSSIAN03.
82. Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648.
83. Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785.
84. Stephens, P. J.; Devlin, F. J.; Chabalowski, C. F.; Frisch, M. J. *J. Phys. Chem.* **1994**, *98*, 11623.
Implementation.
85. Leininger, T.; Nicklass, A.; Stoll, H.; Dolg, M.; Schwerdtfeger, P. *J. Chem. Phys.* **1996**, *105*, 1052.
SDD-basis set.
86. Nifantiev, E. E.; Sorokina, S. F.; Borisenko, A. A.; Zavalishin, A. I.; Vorobjeva, L. A. *Tetrahedron* **1981**, *37*, 3183.
For syntheses of P-H phosphonites (i.e. (RO)₂P-H).
87. Centofanti, L. F. *Inorg. Chem.* **1973**, *12*, 1131.
88. Ostermeier, M.; Prieß, J.; Helmchen, G. *Angew. Chem.* **2002**, *114*, 625.
For applications of hydridophosphanes, i.e.: R₂P-H, in enantioselective catalyses.
89. Nikonov, G. I.; Grishin, Y. K.; Lemenovskii, D. A.; Kazennova, B. N.; Kuzmina, L. G.; Howard, J. A. K. *J. Organomet. Chem.* **1997**, *547*, 183.

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the Beilstein Journal of Organic Chemistry terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at:
[doi:10.1186/1860-5397-1-6](https://doi.org/10.1186/1860-5397-1-6)